

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

FINAL REPORT

TO

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

MISSISSIPPI TEST FACILITY


Contract Number-
NAS -9- 14501

(NASA-CR-147585) EFFECTS OF ENVIRONMENTAL
CHANGES ON MARSH VEGETATION WITH SPECIAL
REFERENCE TO SALINITY Final Report, 1 Feb.
1975 - 31 Jan. 1976 (Tulane Univ.) 112 p HC
\$5.50

N76-22616

Unclas
26787

CSCI 08J G3/42



Alfred E. Smalley
Leonard B. Thien

TULANE UNIVERSITY
New Orleans, Louisiana 70118

February 1, 1976

INTRODUCTION

The following is the final report on the NASA grant "Effects of Environmental Changes on Marsh Vegetation with Special Reference to Salinity." The period of study began on February 1, 1975 and was terminated January 31, 1976.

LITERATURE REVIEW OF MARSHES

The word "marsh" connotes low wet land, often treeless that is periodically inundated and characterized by a variety of grasses, sedges, rushes, etc. Despite the general appearance of marshes, ecologists recognize many different types of marsh communities, i.e., salt marsh, cattail marsh, freshwater marsh, etc.

The following literature review is concerned primarily with brackish and salt marshes located along the eastern coast of North America and the Gulf Coast. The review concentrates upon the vegetation of the marshes particularly in regard to distribution, composition, succession and productivity although other aspects are also reviewed. Special efforts were made to include major works concerned with the Louisiana and Mississippi coastal marshes.

Distribution

Within brackish and salt marshes the phanerogamic vegetation is composed primarily of grasses, rushes and succulent dicotyledons. Such genera as Spartina, Distichlis, Juncus, and Salicornia are encountered repeatedly in widely separated areas (Adams, 1963; Chapman, 1960). Monospecific communities of Spartina sp., Juncus roemerianus and of Distichlis sp. frequently dominate large areas depending upon environmental factors.

The salt marshes begin as grassy arctic marshes in the far north beginning in the northern reaches of Ellesmere Island within the Arctic

circle, on Baffin Island, along the northern shore of Quebec on Hudson Bay and along the upper reaches of Labrador, followed by marshes tucked in coves and bays along the glaciated coast of the Canadian Maritimes and New England coasts. The salt marshes cover significant areas along the central coast of Nova Scotia. Extensive marshes lie behind the barrier beaches of the mid-Atlantic coast. At the southern limit of the salt marshes in Florida, mangrove trees replace the marsh (Teal and Teal, 1969). Salt marshes are found in their greatest abundance from Albermarle Sound on the coast of North Carolina south to the northern coast of Florida and in Louisiana (Teal and Teal, 1969; Martin, Hotchkiss, Uhler and Bourn, 1953; Thorne, 1954; Linton, 1968; Uphof, 1941).

In the South Atlantic and Gulf states there are approximately 5,600,000 acres (22,662 square kilometers) and of this area 3,381,500 acres (13,683 square kilometers) are located in Louisiana (Penfound and Hathaway, 1938; Griffitts, 1928). Louisiana has nearly twelve times as much salt marsh as New Jersey and almost half the total salt marsh area of the Atlantic and Gulf Coasts (Penfound and Hathaway, 1938). Cibula (1972) states that southeastern Louisiana alone possesses 1,750,000 acres (7,123 square kilometers) of fresh water marsh and about 2,500,000 acres (10,117 square kilometers) of salt marsh. The salt marsh vegetation on the northern portion of the Gulf of Mexico extends in a discontinuous arc from Galveston, Texas to just south of St. Petersburg, Florida (Eleuterius, 1973). In addition to the vast quantities of

salt marsh in Louisiana there are approximately 315,000 acres (1,277 square kilometers) in Texas, 26,500 acres (109 sq. km.) in Mississippi, 34,000 acres (138 sq. km.) in Alabama and 680,000 acres (2,873 sq. km.) in Florida (Thorne, 1954; Griffitts, 1928).

Vegetation

The early work on marshes in the United States was done on the Atlantic Coast, especially the northern part of Kearney (1900, 1901), Harshberger (1900, 1909), Johnson and York (1915), Taylor (1939), Conrad (1935), Rudolphs (1926), and Chapman (1940a, 1940b). The southern portion was studied by Wells (1928), Davis (1943), Jackson (1952), Bourdeau and Adams (1956), Bourn and Cottom (1950), Kerwin (1966), Reed (1947), Hinde (1954), McCormick, Grant and Patrick (1970) and Kerwin and Pedigo (1971).

In a paper on the vegetation of Wequetequock-Pawcatuck tidal marsh in Connecticut, Miller and Egler (1950) list botanical literature by state for the Atlantic Coast of North America. Although incomplete, the list is useful. Uphof's (1941) world review of the halophyte problem includes a bibliography of 363 titles.

In the tidal marshes of the north Spartina alterniflora borders the bay, ditches and estuaries. Panicum virgatum and its associates border the uplands. Large colonies of Juncus sp. are found with stretches of Spartina patens and beds of Distichlis sp. distributed about. Forbs such as Plantago sp. and Limonium sp. are scattered

(Miller and Egler, 1950). Thus a typical cross-section of vegetation from bay to upland would be as follows: Spartina alterniflora-lower border, Spartina patens-lower slope, Juncus sp.-upper slope, Panicum virgatum-upper border, Iva sp. and other shrubs-upland shrub border, and Quercus sp. and other forest trees-upland forest. The transition between S. alterniflora and S. patens is marked by the normal high tide zone. Exceptional high tide zone marks the border between Panicum virgatum and the shrub border (Miller and Egler, 1950). A list of plants for the Wequetequock-Pawcatuck Tidal Marsh are presented with frequencies by Miller and Egler (1950). In unit areas along transects on the high marsh at the western end of Barnstable Marsh, the angiosperm communities were dominant in the following proportions (Blum, 1968).

Monospecific Dominance

| | |
|------------------------------|-------|
| <u>Spartina patens</u> | 39.6% |
| <u>Spartina alterniflora</u> | 32.2% |
| <u>Distichlis spicata</u> | 12.0% |
| <u>Juncus gerardi</u> | 1.1% |

Heterospecific Dominance

| | |
|--|-------|
| <u>Spartina alterniflora</u> | 10.0% |
| <u>Spartina patens</u> - <u>Distichlis spicata</u> | 1.6% |
| <u>S. alterniflora</u> - <u>D. spicata</u> | 0.6% |

In the Great Marshes, Barnstable, Mass., Redfield (1972) traces the development of a typical New England marsh and maintains it takes

from 500-1,000 years or more for high marsh to develop. Nixon and Oviatt (1973) discuss the vegetation and other factors of Bissel Cove on Rhode Island and give lists of plants as well as animals with their interaction.

Succession

In the northern marshes (New England) the general succession sequence is as follows (Chapman, 1940a): primary colonist - Spartina alterniflora - (Spartinetum alterniflora) - Spartina patens - Juncetum gerardi - freshwater. Most workers would consider that the Spartina patens community should be subdivided into a Spartina community and a Distichlis spicata community (Chapman, 1940a; Flowers, 1973).

On Saugus Marsh (near Boston) Chapman (1940a, 1940b) recognized the following consocieties:

| | |
|---|----------------|
| <u>Spartinetum alterniflora glabrae</u> | Depth 2.5 m |
| <u>Spartinetum patentis</u> | 1.5 |
| <u>Spartinetum-Distichlidetum</u> | .4 |
| <u>Juncetum-Gerardii</u> | .4 |
| <u>Spartinetum pectinatae</u> | .4 |
| <u>Scirpetum</u> with Phragmites | - |

Chapman (1960) presents a series of diagrams of plant succession at various locations in northern marshes. The Spartinetum alterniflora glabrae is a "sedge association" (Conrad, 1935; Ganong, 1903) and is

recognized from Nova Scotia through the northern shores of the Gulf of Mexico. The next shoreward community is either the Spartinetum patentis, the Distichlidetum spicatae or Salicornietum ambiguae (Conrad, 1935). Rather than use the above association names it is perhaps best to simply list species for each successional zone. It should be noted that Spartina alterniflora is the first species to appear starting with bare mud in saline areas on the coast (Chapman, 1960).

Southern marshes

Species associations in the North Carolina salt marshes are given by Adams (1963).

Low Marsh

Spartina alterniflora
S. alterniflora-S. perennis, Limonium carolinianum
Juncus roemerianus

High Marsh

Aster tenuifolius, Distichlis spicata,
Fimbristylis castanea, Borrichia frutescens
and Spartina patens.

The primary colonist is Spartina alterniflora. It may contain Salicornia perennis or Limonium carolinianum and these plants reach their best development at about mean sea-level. At higher levels Juncus roemerianus is the dominant.

South Atlantic Coast

Successional Series for the south Atlantic Coast is as follows:

Spartina alterniflora - Spartina sp. - Salicornia sp. - Juncus sp. -
Fimbristylis sp. - Baccharis sp. - Kosteletzkya sp. Various diagrams
showing succession are given by Wells (1928). Arranged according to
depth the species are associated in the following manner (Chapman,
1960).

Deeper Water:

Spartina sp. and Salicornia sp.

Middle Zone: (shallow water)

Distichlis spicata
Borrichia futescens
Juncus sp.
Lythrum lineare
Scirpus olneyi

Inner Zone:

Baccharis - Kosteletzkya

The salt marshes of Florida are slightly different in that there
is a transition to mangrove swamp. In the northern portion of the
state, the successional relationships are similar to those from Virginia
and the Carolinas. In the southwest, the salt marsh commonly develops
behind the mangrove swamp or enclaves with the mangrove (Chapman, 1960).
The community succeeding the mangrove is often dominated by Batis maritima
and at higher levels is associated with Salicornia ambigua. A Distichlis
sp. - Spartina patens community occurs at still higher levels followed
by a Juncus roemerianus - Spartina cynosuroides - Spartina bakeri
community (Chapman, 1960). Jackson (1952) analysed the covital vegetation

within the St. Marks National Wildlife Refuge in Wakulla County, Florida.

He lists the vegetation of various zones (A - D) in order of abundance.

Zone A (only the edge of this zone is inundated)

Fimbristylis castanea
Spartina patens
Spartina spartinae
Limonium carolinianum
Juncus roemerianus
Salicornia ambigua
Distichlis spicata
Batis maritima
Seutera nitida

Zone B - S ("salt-zone")

Salicornia ambigua

Zone C

Distichlis spicata
Spartina cynosuroides
Salicornia ambigua
Batis maritima
Borrichia frutescens
Juncus roemerianus
Limonium carolinianum

Zone D

Juncus roemerianus
Limonium carolinianum
Batis maritima

Louisiana & Mississippi

Eleuterius (1973) notes that very little work has been done on the marshes associated with the Gulf of Mexico until recent years. The cornerstone work on the Louisiana marshes (Gulf Coast marshes in general) was by Penfound and Hathaway (1938). Seven transects were laid out in

the marshes south of New Orleans from fresh to salt water involving eleven distinct community types. The paper gives extensive lists of plants found within these various communities.

The marshes on the coastline of the northern Gulf of Mexico vary considerably. West of the Mississippi River to the 98th meridian in Texas, Spartina spartinae and Sporobolus virginicus are the dominant species (Tharp, 1926). A vegetational map of the marshes of Louisiana by Chabreck et al. (1968) shows little saline marsh west of the 91st meridian but large expanses of brackish marsh are evident. The works of Day (1959), Pullen (1960, 1962), Childress (1960) should be consulted for lists of plants and maps of marshes along the Texas coast. Hoese (1967) also maintains that there is little or no salt marsh west of Vermilion Bay, Louisiana and that little Spartina alterniflora marsh is present in Mississippi with the Apalachicola Bay marshes poorly defined.

In Mississippi, Juncus roemerianus is the dominant plant species in the marshes (Eleuterius, 1973). Line transect data revealed that this species composed 57.8% of the marsh population in April, 42.1% in June and 43.7% in August. The marsh was not as diverse in species in April (25 species) as it was in June (43 species) and August (40 species). J. roemerianus composed 45.3% of the total plant population for the entire growing season (Eleuterius, 1973).

The twelve most abundant plant species in the marshes of Mississippi are as follows: J. roemerianus 42%, Spartina patens 8%, Spartina alterniflora 7%, Spartina cynosuroides 6.5%, Cladium jamaicense 3%, Scirpus

validus 2.5%, Distichlis spicata 2%, Osmunda regalis 1.5%, Fimbristylis spadicea 2%, Phragmites communis 1.5%, and Boltonia asteroides 1% (Eleuterius, 1973).

In Mississippi the primary difference between the brackish and saline marshes is the reduction in the abundance of Spartina alterniflora and an increase in brackish and fresh water plant species (Eleuterius, 1973).

In the saline marshes two species predominate: Juncus roemerianus and Spartina alterniflora. Brackish water species, such as Spartina cynosuroides, Spartina patens and Scirpus olneyi are found intermixed with J. roemerianus. S. alterniflora always occurs in pure stands (Eleuterius, 1973).

Distichlis spicata, "salt grass" and several succulent plants as Salicornia bigelovii, Suaeda linearis and Batis maritima grow in the "salt flats" area. J. roemerianus is always associated with these plants (Eleuterius, 1973).

Scirpus olneyi, S. robustus and Spartina patens occur as zones near the periphery of the marsh.

In the intermediate marsh (overlap between brackish and freshwater) the upper limit of Juncus roemerianus is reached. Phragmites communis becomes very common in this area (Eleuterius, 1973).

Louisiana

The Gulf Coast marshes of Louisiana range in elevation from minus 6.096 dm. (2 feet) to 6.096 dm. above sea level (excluding chenieres

and natural levees). Most of the marshes in Louisiana are on land created by the Mississippi River through thousands of years. The Mississippi River has inhabited seven courses in the past 7,200 years (Juneau, 1975). About 2,800 years ago the river shifted eastward to occupy the St. Bernard course in southeastern Louisiana. While occupying the St. Bernard course, the Mississippi River developed a vast delta extending from the general vicinity of Barataria Bay out into the Gulf beyond the present position of the Chandeleur Island group (National Shoreline Study, 1971). Then about 1,200 years ago, the Mississippi River again shifted to the west. Today the Breton and Chandeleur Island groups represent a late stage in deltaic distribution resulting from subsidence behind the old shoreline (National Shoreline Study, 1971). The present delta began building about 400 years ago (Russell, 1936; Coleman, 1966).

In Louisiana, O'Neil (1949) classifies the marsh into Delta marsh, Sub-delta marsh and Prairie marsh. The Delta marshes consist of approximately 300,000 acres (1192 sq. km.) centered around the Mississippi River into the Gulf of Mexico (present day outlet).

The Sub-delta marshes consist of approximately 2,940,000 acres (11,895 sq. km.) which start at Baptiste Collette and Grand Pass, Plaquemines Parish on the southeast, and continue in a meandering line from Cow Island to the east of Cheniere au Tigre in Vermilion Parish on the west. The Sub-delta marshes are the ancient deltas of the Mississippi River (O'Neil, 1949).

The prairie marshes are located in Vermilion and Cameron Parishes and consist of approximately 760,000 acres (2,670 sq. km.) according to O'Neil (1949).

Other papers dealing with the vegetation of the Gulf marshes are by Curl (1959), Kurz and Wagoner (1957) on the Florida marshes. The Mississippi and Louisiana marshes have been described by Lloyd and Tracy (1901), Lowe (1921), Penfound and O'Neil (1934), Penfound and Hathaway (1938), Penfound (1952), Viosca (1928) and Mohr (1901), Lemaire (1961), Egglar (1961), Eichhorn, and Duice (1969).

In recent years there has been a flourish of activity in analyzing the vegetation of the marshes of Louisiana and Mississippi with works by Chabreck (1970), Uhler and Hotchkiss (1968), Gabriel and de la Cruz (1974), Shiflet (1963), Linton (1968), Palmisano (1970), Juneau (1975) and Woodhouse, Seneca and Broome (1974). A vegetation map of the Louisiana coastal marshes showing saline, fresh, brackish and intermediate vegetation has been constructed by Chabreck, Joanen and Palmisano (1968). Also a phytogeographic and ecologic relationships of the flora of Breton Island has been written by Gould and Ewan (1975). For notes on Louisiana in general with keys to biting flies consult Tidwell (1973).

Succession

Few studies of succession have been made in the Gulf coast area. However, Chapman (1960) maintains that the general succession within

the marshes is not materially different from that of the New England marshes except at the higher levels, or where, in Florida and Texas, there is a transition to the mangrove area (also Louisiana). Chapman (1960) gives a few diagrams of succession in the Gulf area.

Penfound and O'Neil (1934) give a successional sequence on Cat Island as follows (for brackish water): *Spartina* Conspecies - *Distichlis* Conspecies - *Juncus* Conspecies.

On Grand Isle, Louisiana, Walker (1939) gives a sequence as follows: *Spartina alterniflora* - *Distichlis* and *Avicennia* - *Baccharis* - *Iva* association.

Factors Affecting Zonation and Distribution in Marshes

In an excellent paper Adams (1963) discusses factors influencing plant zonation. Chapman (1940a, 1940b) thought inundation was the important factor while Miller and Egler (1950) considered the present day distribution of plants due to past environmental changes that could not be interpreted today. Johnson and York (1915) believed salt marsh species to be distributed according to submergence-to-emergence ratios. Reed (1947) working in North Carolina considered that inundation, salinity and poor drainage were important factors at lower levels of the marsh whereas competition with other angiosperms on the upper periphery was the main factor.

Adams (1963) studied the effects of elevation, soil texture and salinity on the distribution of *Spartina patens*, *Juncus roemerianus*, and the tall, medium and short growth forms of *Spartina alterniflora*.

A micro-relief gradient was found to be significant in delimiting the lower boundary of Spartina patens. Soil texture was similar for J. roemerianus and S. alterniflora. Salinity of the soil at low tide increased landward to a value more than twice as high in the short growth form of S. alterniflora and Salicornia perennis and then decreased to a low level in the S. patens zone. Adams (1963) concludes that tide level elevation influences are the primary factors controlling the distribution of salt marsh species. Most salt marsh species exhibit reduced growth and fertility with increasing salinity-salt concentrations of about 7% NaCl (twice sea strength) prohibit establishment and survival of all species.

Apparently S. alterniflora is restricted to the low marsh because of its moderate salinity and high iron requirements (Adams, 1963). If grown in low iron or fresh water it becomes chlorotic whereas Distichlis spicata and Spartina patens (competitors) do not display this quality. Thus Distichlis spicata and S. alterniflora are salt obligates and S. patens a facultative halophyte (Adams, 1963).

Babcock (1967) in a study of wiregrass (Spartina patens) and saltmarsh grass (Distichlis spicata) in Louisiana investigated the effects of salinity and water depth. It was found that both species grew best in salinity of 0.50 - 2.50‰. At salinities lower than this both species were greatly reduced in number and replaced by Scirpus olneyi. It was thought that water level was a very important limiting factor. As water depths exceeded .3048 m (1 foot), the densities of

wiregrass and saltmarsh grass decreased with young sprouts more sensitive to flooding than older shoots.

Chapman (1960) maintains that germination of many salt marsh species is dependent on a reduction in the soil surface salinity. Germination tests with seeds of S. alterniflora indicate decreased germination rates with increasing salinity (Mooring, Cooper, and Seneca, 1971). The height forms of S. alterniflora apparently represent different ecotypes that result from exposure to environments differing in salinity (Mooring et al., 1971), (Woodhouse, Seneca, and Broome, 1974), (Shea, Warren, and Niering, 1975). For an excellent description of Spartina species see Mobberly (1956). This taxonomic treatment also provides keys to species as well as distribution maps for each species.

In Louisiana, Penfound and Hathaway (1938) emphasize that the transition from one community to another is conditioned by a change in elevation of as little as 7.62 cm since four communities may occur on a slope having a fall of less than 3.048 dm. Penfound and Hathaway (1938) maintain that water level and salinity of the soil water are the most important habitat factors influencing the plants in the Louisiana marshes. In the probable order of salt tolerance the major marsh dominants in Louisiana are (Penfound and Hathaway, 1938):

| | |
|---------------------------------------|--------------------------|
| <u>Mariscus (Caldium) jamaicensis</u> | %Salt Tolerance 0-0.2 |
| <u>Scirpus californicus</u> | 0-1.13 |

| | |
|------------------------------|-----------|
| <u>Typha latifolia</u> | 0-1.13 |
| <u>Scirpus olneyi</u> | 0.55-1.68 |
| <u>Typha angustifolia</u> | 0-1.68 |
| <u>Spartina patens</u> | 0.12-3.91 |
| <u>Juncus roemerianus</u> | 0.12-4.43 |
| <u>Spartina alterniflora</u> | 0.55-4.97 |
| <u>Distichlis spicata</u> | 0.45-4.97 |
| <u>Avicennia nitida</u> | 3.68-4.97 |

Isohaline map of Louisiana coastal marshes see Chabreck and Palmisano (1968). A map of isohalines for the Breton Sound area (specifically the Mississippi River Gulf Outlet Canal area) is presented by Rounsefell (1964). The vegetation map of Chabreck, Joanen and Palmisano (1968) list the following common plants for the marshes of Louisiana:

Saline

S. alterniflora
Salicornia sp.
Juncus roemerianus
Batis maritima
Avicennia nitida
Distichlis spicata

Brackish

S. patens
Scirpus olneyi

Intermediate

Vigna repens

Scirpus californicus

Cladium jamaicense

Fresh

Panicum hemitomon

Alternanthera philoxeroides

Hurricanes

Hurricane damage to any area is largely dependent on the physical features of that area. Usually hurricanes produce an increase in the amount of open areas, effect translocation of vegetation inland, cause deposits of vegetational debris along levees, reduce the density of dominant plant species, plus raise the salinity, particularly of fresh water marshes.

The effect of hurricane Audrey on Marsh Island, Louisiana, was studied by Harris and Chadreck (1958). Marsh Island, a saltmeadow-cordgrass (Spartina patens) and Olney's three square (Scirpus olneyi) marsh, became flooded with about 2.5 m of water. The normal tide water at Marsh Island is brackish because fresh water from the Atchafalaya Basin bathes the island shores. Salinities of soil water measured during 1955 and 1956 ranged from 7 to 29 per cent sea-water, while measurements taken after the hurricane were 8 to 20 per cent of sea water. If salinity of the water is altered by the hurricane, it quickly returns to its normal range.

A change in the dominant vegetation of the brackish marsh on Marsh Island was noticed after the hurricane. In 1956 Spartina patens

comprised 99% of the brackish marsh and Scirpus olneyi 1%. In 1958 S. patens comprised 90.6 and S. olneyi 9.4% (Harris and Chabreck, 1958).

Chamberlain (1959) studied the influence of hurricane Audrey on the Rockefeller Wildlife Refuge, Louisiana, an area about 45 km west of Marsh Island. The tidal surge was at least 2.5 m in this area.

Natural beaches, formed of sand and shells in equilibrium with the tide and wave action, separates the Gulf of Mexico from the salt-water marsh. These beaches, characterized by a relatively low profile, underwent a geomorphic change due first to the rising tide water and wave action, then to the recession of the tidal surge. Storm waves broke over the submerged crest causing mass transport of and spreading of shell-sand material. This resulted in the rear margin of the beach being moved approximately 100 m. The main change in the beach habitat was in the distribution of the endemic flora. There was a general increase in soil salinity, but in the salt marsh and brackish zone this increase has very little effect. In the fresh water zone, however, the increase was disastrous. Where dense stands of Cladium jamaicense was dominant the intrusion of salt water was severe. Pre-hurricane growth of this species in the Sweet Lake region was such that travel through this area was almost impossible. After the hurricane the field resembled a wet prairie. C. jamaicense has been in a steady decline in

the southwest area of Louisiana largely due to salt-water intrusion and impeded marsh drainage.

Two large masses of clay were deposited over the beach by the hurricane's tidal surge and these "mud arcs" have become permanent features of the shoreline. Four weeks after the hurricane the mud was very fluid, six months later the mud surface dried, becoming cracked. Fourteen months after the formation of the mud arcs a study was made to determine the type and density of the vegetation. The vegetation-encroachment, on the peripheral of the mud arc, is a typical salt marsh with Spartina alterniflora, Distichlis spicata, Borrichia frutescens and Batis maritima. It was estimated that it would take approximately two years for the arcs to become completely covered with vegetation.

Chabreck and Palmisano (1973) studied the effect of hurricane Camille on the marshes of the Mississippi River delta. Immediately following the hurricane the salinity in the marsh along the natural levee increased from .92 ppt to 3.61 ppt. But one year after the hurricane the salinity was raised from .10 ppt to .26 ppt but receded to .11 ppt within one year. Alternanthera philoxeroides seemed to be the species affected by the increase in salinity. This species was absent in the ponds and lakes after the hurricane and had not reappeared one year later.

In general in the salt marsh and brackish zones recovery after a hurricane can be complete after one or two uninterrupted growing seasons.

Productivity

Estuaries in general and salt marshes in particular are unusually productive places (Teal and Teal, 1969). None of the common agriculture, except possibly rice and sugarcane production, comes close to producing as much potential animal food as do the salt marshes. The agricultural crops which approach this high figure are fertilized and maintained by man at great expense (Teal and Teal, 1969). There are many reasons to account for the productivity of marshes (particularly salt marshes). Only a small percentage (less than 10%) of the organic material produced by the marsh plants is actually grazed by marsh herbivores (de la Cruz, 1973; Keefe, 1972). The bulk of the plant materials dies (annually for most species) and falls to the surface of the mud where it may decompose (Teal and Teal, 1969). Protein enrichment due to bacteria occurs shortly thereafter.

Another reason for the richness of marshes (salt) is that there is a continuous mixing of nutrients (and water) in the marshes. These nutrients are turned over at a relatively high rate in some cases. Pomeroy, Johannes, Odum, and Roffman (1969) found that the sediment in the well-established coastal Spartina alterniflora marsh contained enough phosphorous for 500 years growth without replenishment. Their studies indicate that Spartina obtains its phosphorous from the soil. The phosphorous in the water provides phosphorous for the sediment for it is absorbed on the surface of the sediment, then combined into the crystal lattice of the clay. From this combination with the clay, it

can still be released as free phosphate available for biological uptake. Thus, the large supply of phosphorous in the sediment assures a continuing source of phosphorous for phytoplankton and mud algae, as well as Spartina.

The high concentration of organic matter in the soil leads to the formation of colloids. The colloids absorb ions necessary for plant growth (Albrecht, 1941; Gorham, 1953). The decomposition of organic matter produces large quantities of carbon dioxide, increasing the acidity of the water and thereby increasing the solubility of iron and manganese (Keefe, 1972). These ions may occur at concentrations that would normally be considered toxic (Robinson, 1930), but marsh plants have adapted to high concentrations. Spartina has an unusually high demand for iron and despite the high concentrations in marsh soil still appears chlorotic in some areas (Teal and Teal, 1969).

The tides not only remove detritus from marshes but also add nutrients (Aurand, 1968). In addition, flowing water is more important than standing water (Schelske and Odum, 1961). Particles rich in nutrients are trapped on the marsh obstructions provided by the plants (Gorham and Pearsall, 1956; Blum, 1969; Ranwell, 1964a), thus increasing the size of the marsh laterally and vertically, while producing soil rich in nutrients for the plants. Soil nutrients decrease in concentration away from the sea (Ranwell, 1964b).

Nutrient levels in the water over freshwater marshes are often high. Buttery, Williams, and Lambert (1965) reported values of 0.17-0.30 mg $\text{PO}_4\text{-P}$ liter⁻¹, 0.81-1.18 mg $\text{NO}_3\text{-N}$ liter⁻¹ and 2.88-1.08 mg

$\text{NH}_3\text{-N}$ liter $^{-1}$ in the water over a Phragmites communis-Glyceria maxima streamside marsh. Phosphorous in the soil water decreased from 100.9-509 mg liter $^{-1}$ at the channel to 9.5-9.3 mg liter $^{-1}$ at the landward side of the marsh.

Another reason for the high productivity of marshes (salt) may be the vertical orientation of the leaves. This orientation reduces intenss heating (Palmer, 1941), exposes the maximum leaf surface to sunlight over the day, and minimizes mutual shading (Jervis, 1964).

de la Cruz (1974) analyzed nine species of marsh plants in the Gulf Coast of Mississippi to determine their productivity (monotypic stands). Three 1 square meter plots were harvested monthly from each of the nine plots. Annual net productivity of the above ground material was estimated from the monthly increases during the season. Caloric values of the plant material were also recorded along with ash-free weight. The values are as follows:

| Species | Annual Net Primary Productivity | |
|------------------------------------|---------------------------------|---------------------|
| | Dry g/m ² | Kcal/m ² |
| <u>Sagittaria lancifolia</u> | 600 | 2468 |
| <u>Phragmites communis</u> | 2330 | 9841 |
| <u>Scirpus robustus</u> | 1056 | 4576 |
| <u>Juncus roemerianus</u> | 1697 | 7558 |
| <u>Spartina cynosuroides</u> | 2190 | 9347 |
| <u>Spartina patens</u> | 1922 | 8464 |
| <u>Distichlis spicata</u> | 1484 | 6020 |
| <u>Spartina alterniflora</u> short | 1089 | 4028 |
| <u>Spartina alterniflora</u> long | 1964 | 8088 |

The production value of $1697 \text{ g/m}^2 \text{ yr}^{-1}$ for Juncus roemerianus is higher than previously reported in the literature (Williams and Murdoch, 1972; Heald, 1969; Stround and Cooper, 1968; Waits, 1967).

The net primary productivity of S. alterniflora short form is higher than previously reported; the tall form productivity is comparable to the values reported for Georgia and Louisiana (de la Cruz, 1973; Keefe, 1972). The values for the other species are comparatively higher than the production values observed for similar and related species from other estuaries. de la Cruz (1974) found the tidal marshes to be generally more productive than the marshes found in the Atlantic Seaboard. Dirby (1971) suggested that the greater productivity in the Gulf marshes may be partly due to the longer growing season.

Keefe (1972) in a major review article gives a summary of biomass and production values for marsh plants. The list of over sixty items covers plants growing in salt, brackish and freshwater growing throughout the Atlantic and Gulf Coasts.

de la Cruz (1973) also provides a list on estimates of annual Net Primary Production shown in Table A.

In the salt marsh, the water flooding the marsh is the source of the nutrients, but there is a net loss of produced material to the estuary, providing food for consumers (Odum and de la Cruz, 1963; Reimold, 1965; Reimold and Daiber, 1970). In addition there is evidence that salt marshes serve as spawning and nursery grounds for the young of a number of economically important marine species (see Hitchcock, 1972 for a review).

TABLE A. Values from de la Cruz, 1973

| Marsh Community | Annual Net Production g/m ² | Geographic Location | Reference |
|------------------------------|--|-----------------------------|---|
| Mixed vegetation | 992-1108 | St. Louis Bay Estuary, Ms. | de la Cruz & Gabriel (Present Research) |
| Mixed vegetation | 1246 | Patuxent Estuary, Md. | Johnson (1970) |
| <u>Spartina alterniflora</u> | 2883 | Sapelo Is., Ga. | Odum & Fanning (1973) |
| <u>Spartina alterniflora</u> | 1150 | Barataria Bay Estuary, La. | Kirby (1971) |
| <u>Spartina alterniflora</u> | 1207 | Patuxent Estuary, Md. | Johnson (1970) |
| <u>Spartina alterniflora</u> | 650 | Beaufort, N.C. | Williams & Murdoch (1969) |
| <u>Spartina alterniflora</u> | 1158 | Sapelo Is., Ga. | Teal (1964) |
| <u>Spartina alterniflora</u> | 2000 | Sapelo Is., Ga. | Schelske & Odum (1961) |
| <u>Spartina alterniflora</u> | 445 | Canary Creek Estuary, Del. | Morgan (1961) |
| <u>Spartina alterniflora</u> | 985 | Sapelo Is., Ga. | Smalley (1959) |
| <u>Juncus roemerianus</u> | 2000 | Ocean Springs, Ms. | Eleuterius (Pers. comm.) |
| <u>Juncus roemerianus</u> | 754 | Cape Lookout, N.C. | Williams & Murdoch (1972) |
| <u>Juncus roemerianus</u> | 849 | Everglades, Fla. | Heald (1969) |
| <u>Juncus roemerianus</u> | 796 | Cape Fear River, N.C. | Stroud & Cooper (1968) |
| <u>Juncus roemerianus</u> | 560 | Cape Fear River, N.C. | Foster (1968) |
| <u>Juncus roemerianus</u> | 895 | Bodie Is., N.C. | Waits (1967) |
| <u>Spartina cynosuroides</u> | 1028 | Altamaha River Estuary, Ga. | Odum & Fanning (1973) |
| <u>Spartina patens</u> | 1296 | Bodie Is., N.C. | Waits (1967) |
| <u>Spartina patens</u> | 993 | Long Is., N.Y. | Harper (1918) |

Decomposition

Table B gives a summary of decomposition rates for various marsh plants. The data represents the fate of plant materials being produced and dying on the marsh. Decomposition rates are expressed as per cent loss of material. The removal of material from the surface of the marsh by water makes the plant material available to the detritus-based consumers (de la Cruz, 1973; de la Cruz and Gabriel, 1974).

Energy Flow

Organisms maintain themselves and reproduce by utilizing energy from the environment. This movement of energy from organism to organism is called energy flow (McNaughton and Wolf, 1973). Analysis of energy flow in ecosystems is a tedious undertaking that takes years of effort.

A summary of energy flow in a Georgia salt marsh has been worked out by Teal (1962). A summary of his findings is given below:

| | |
|---------------------------------------|----------------------------------|
| Input as light | 6000,000 kcal/m ² /yr |
| Loss in photosynthesis | 563,620 or 93.9% |
| Gross production | 36,380 or 6.1% of light |
| Producer respiration | 28,175 or 77% of gross prod. |
| Net production | 8,205 kcal/m ² /yr |
| Bacterial respiration | 3.890 or 47% of net prod. |
| Primary consumer respiration | 596 or 7% of net prod. |
| Secondary consumer respiration | 48 or 0.6% of net prod. |
| Total energy dissipation by consumers | 4,534 or 55% of net prod. |
| Export | 3,671 or 45% of net prod. |

TABLE B. Decomposition rates of various plant species expressed as % loss of material after one year. From de la Cruz, 1973.

| Species | Decomposition Rate (%/yr) | Locality | Methods | Reference |
|---------------------------|---------------------------|--------------------------|------------------------------------|---|
| <u>Juncus roemerianus</u> | 40 | Bay St. Louis, Ms. | Nylon net bags, 5.0 mm mesh | de la Cruz and Gabriel (1973) |
| <u>Juncus roemerianus</u> | 46 | N. of Cape Lookout, N.C. | Dissappearance rate, tidal removal | Williams and Murdoch (1972) |
| <u>Juncus roemerianus</u> | 36 | Everglades, Fla. | Nylon net bags, 1.5 mm mesh | Heald (1969) |
| <u>Juncus roemerianus</u> | 47 | Bodie Is., N.C. | Nylon net bags, 1.0 mm mesh | Waits (1967) |
| <u>Juncus roemerianus</u> | 35 | Sapelo Is., Ga. | Nylon net bags, 2.5 mm mesh | de la Cruz (1965) |
| <u>Juncus squarrosus</u> | 27 | Lancashire, England | Nylon net bags, 10 mm mesh | Latter and Cragg (1967) |
| <u>Scirpus americanus</u> | 60 | Bay St. Louis, Ms. | Nylon net bags, 5.0 mm mesh | de la Cruz and Gabriel (Present Research) |
| <u>Distichlis spicata</u> | 38 | Bay St. Louis, Ms. | Nylon net bags, 5.0 mm mesh | de la Cruz and Gabriel (Present Research) |
| <u>Distichlis spicata</u> | 53 | Sapelo Is., Ga. | Nylon net bags, 2.5 mm mesh | de la Cruz (1965) |
| <u>Salicornia sp.</u> | 94 | Sapelo Is., Ga. | Nylon net bags, 2.5 mm mesh | de la Cruz |
| Sawgrass leaves | 45 | Everglades, Fla. | Nylon net bags, 2.5 mm mesh | Heald (1969) |
| Red mangrove leaves | 60 | Everglades, Fla. | Nylon net bags, 2.5 mm mesh | Heald (1969) |
| White mangrove leaves | 38 | Everglades, Fla. | Nylon net bags, 2.5 mm mesh | Heald (1969) |

In the marsh analyzed Spartina accounts for approximately three-fourths of the primary production, while algae on the mud accounts for the rest. The model shows that only 1.5% of the sunlight is transformed into energy available to consumers. In other marshes, even less incident radiation is converted to organic matter. In a Spartina alterniflora - Juncus roemerianus marsh in North Carolina Stroud and Cooper (1969) found that only 0.2% of the incident light was converted to net primary production. Bray (1962) found 0.6% of the incident light was utilized in a Typha angustifolia - latifolia hybrid marsh in Minnesota.

In Teal's study (1962), 45% of net production was exported to the surrounding waters. This export would be available to the estuarine consumers. This is quite high compared to similar ecosystems. Nixon and Oviatt (1973) estimated 10 to 30% was exported in a salt marsh in Rhode Island (also gives energy flow diagrams).

Energy flow studies for the Gulf Coast similar to Teal's (1962) and Nixon and Oviatt (1973) for New England area have not been conducted.

SITES

The study areas are located in southeastern Louisiana along Bayou Terre Aux Boeufs and adjoining bays. See the attached map for location of the following study sites.

| Site No. | Name | Location | | | |
|----------|----------------|----------|----|----|----|
| 1 | Petain Lagoon | 89 | 42 | 29 | 45 |
| 2 | Bayou Gaudet | 89 | 42 | 29 | 44 |
| 3 | Bottle Lagoon | 89 | 39 | 29 | 42 |
| 4 | Dead Duck Pass | 89 | 37 | 29 | 41 |
| 5 | Drum Bay | 89 | 34 | 29 | 39 |


Petain Lagoon (1) is a mixed stand of vegetation with species of Spartina alterniflora, Spartina patens, and Spartina cynosuroides with areas of Distichlis spicata and Juncus roemerianus.

Bayou Gaudet (2) consists of an unburned area of Spartina patens and a burned area of Distichlis spicata with very sharp lines between the two species. The margin of this area has many species of halophytes.

Bottle Lagoon (3) is intermediate in the gradient of study plots. Plants within the site include Spartina alterniflora, Spartina patens, Distichlis spicata, and Juncus roemerianus with additional minor species.


Dead Duck Pass (4) study area is composed primarily of Juncus roemerianus with populations of Spartina alterniflora and Distichlis spicata scattered about the site.

Drum Bay (5) is composed almost entirely of Spartina alterniflora with succulent halophytes on the margins including mangrove (Avicennia sp.).



8143 IV SW
(LAKE ELOI 1240)

| | | | | |
|--------|-----|--------|-----|----|
| DO FEE | 756 | R 17 E | 757 | 89 |
|--------|-----|--------|-----|----|



The study area for this report is located in the Sub-delta marsh area of Louisiana (St. Bernard). These marshes are delta formations that have been inactive for possibly 200-500 years in the eastern areas. Their geographic position places them a considerable distance from the Tremendous amounts of fresh water poured into the Gulf of Mexico by the Mississippi River. The mineral deposits are generally of marine origin and saline in nature (O'Neil, 1949). Most of the Sub-delta area is subject to daily tide action. Continual subsidence of the Sub-delta marshes is evident almost anywhere, i.e., the abandoned sea-rims of the Chandeleur Islands. The Breton and Chandeleur Islands have retreated westward at about 13.7 ft/yr (ca. 65 m) between 1812 and 1954. Between 1807 and 1939, the land has been reduced from 721 square miles (1,867 sq. km.) to about 617 square miles (1,598 sq. km.) due to retreat of shoreline (National Shoreline Study, 1971) and enlargement of ponds, bays within the marshlands (O'Neil, 1949). Further evidence are the drowned marshes of Breton Sound (O'Neil, 1949), plus the ghost cypress swamps that flank practically all the bayous near their midway areas, the trees having been destroyed by the intrusion of salt water. Dead live oaks (Quercus virginiana) are present on the crests of natural levees and the vegetation replacing the oaks are the brackish low-ridge species primarily.

Spartina cynosuroides, Baccharis halimifolia and Iva frutescens (O'Neil, 1949).

In the Bayou Terre Aux Boeuf area (location of study plot) the islands are dominated by mangrove, Avicennia nitida. The first substantial marsh in this area is a crest of matted roots and mineral soils 12-15 inches (3.48 dc.) in depth. Under this is an organic peat about 6 m in depth (O'Neil, 1949). About midway on Bayou Terre Aux Boeuf until the direct influence of the present-day river's natural levee Scirpus olneyi dominates. This species has been cultivated for years by annual burning making it easier to hunt animals and travel (O'Neil, 1949). This practice has been used for over 100 years in the area.

The area in general has been greatly disturbed. Hunters and trappers sometimes cut small ditches or drag their piroques over certain spots. Tidal action would in a period of years cut a canal 6 to 10 m wide and 2-4 m deep. The natural bayou levees and lake shores are built of organic material. The entire area is shredded with bayous and dotted with lakes and ponds (O'Neil, 1949).

The marsh in the study area contains Spartina alterniflora, Spartina patens, Spartina cynosuroides, Distichlis spicata, Sporobolus sp. and Fimbristylis sp. (primary species). The marshes of Louisiana are very complex in regard to zonation unlike many of the Atlantic Coast marshes, particularly those found in the New England area. In the Gulf, Spartina alterniflora reaches its lower limit at mean sea level. In the brackish marshes the dominant community is a Spartina patens - Distichlis spicata - Juncus roemerianus complex, but as

the Spartina is less tolerant of salt than the other two species it disappears in the more saline areas. The free soil water in the marshes is more saline than the water in neighboring lagoons and it is thought that the high salinity of the soil water must be maintained by surface evaporation and transpiration (Chapman, 1960).

METHODS

Within each site a 200 meter square study area was delimited and the vegetation analyzed within the square. Two procedures were used to gather vegetation data from these areas. The first method, clipped quadrat (Cox, 1972), was used at all sites. At each site a number of quarter square meter quadrats were clipped. The samples were separated and counted per species. The following measurements were then determined:

$$\text{density} = \frac{\text{number of individuals}}{\text{area sampled}}$$

$$\text{relative density} = \frac{\text{density for a species}}{\text{total density for all species}} \times 100$$

$$\text{frequency} = \frac{\text{number of plots in which species occurs}}{\text{total number of plots sampled}}$$

$$\text{relative frequency} = \frac{\text{frequency value for a species}}{\text{total of frequency values for all species}} \times 100$$

$$\text{importance value} = \text{relative density} + \text{relative frequency} \\ (\text{per species})$$

At those sites in which the vegetation was not dominated by one species, another procedure was also employed. The line transect or line intercept (Canfield, 1941) technique was used with modifications to facilitate its application in the salt marsh. In this study the line consisted of a 4 meter wooden pole subdivided into 1 meter intervals. This pole was run at ground level and the number of individuals per species per interval was recorded. With these values and the average stem diameter values for each species as measured from the

clipped samples, the following measurements were then determined:

$$\text{frequency} = \frac{\text{intervals in which species occurs}}{\text{total number of transect intervals}}$$

$$\text{relative frequency} = \frac{\text{frequency value for a species}}{\text{total of frequency values for all species}} \times 100$$

$$\text{density} = \frac{\text{number of stems per species}}{\text{average stem diameter}} \times \left(\frac{10^3}{\text{number of intervals}} \right)$$

$$\text{relative density} = \frac{\text{density value for a species}}{\text{total density for all species}} \times 100$$

$$\text{dominance} = \frac{\text{number of stems per species (average stem diameter)}}{\text{total transect length}}$$

$$\text{relative dominance} = \frac{\text{dominance value for a species}}{\text{total dominance for all species}} \times 100$$

$$\text{importance value} = \text{relative frequency} + \text{relative density} + \text{relative (per species) dominance}$$

The importance values are seen to be on two scales, 200 for the clipped quadrat method and 300 for the line transect method.

For this study's work in the field the two methods were used in the following manner. At a mixed site a random point of entry along the shoreline was chosen. Once past the shoreline vegetation a quadrat sample was taken. From that point a series of 4 lines was run moving inland. From there 20 paces were traversed and then another quadrat sample was taken. This procedure was followed until the sample area had been crossed. The final sample taken was always a clipped quadrat. (See diagram 1 for example.)




Diagram 1. Sampling method in the field.

The use of these two methods permitted a more detailed analysis of a community's vegetation. Also possible was the corroboration of two unrelated methods for the discernment of any changes noted in the community's structure over time. It is to these ends that these methods were applied.

Salinity measurements were taken concurrently with the vegetation sampling. Both water and soil water were measured when possible. The salinity was measured using a YSI SCT meter.

RESULTS

The following twenty-five tables gives density, relative density, relative frequency, relative dominance and importance values for each site for the period of study.

The following list gives tables for specific study sites:

| | |
|-------------|----------------|
| Table 1-6 | Petain Lagoon |
| Table 7-12 | Bayou Gaudet |
| Table 13-18 | Bottle Lagoon |
| Table 19-24 | Dead Duck Pass |
| Table 25 | Drum Bay |

Salinity values for the various sites are given in Table 26.

In addition, Figure 1 and 2 give density values per week of the year for Bayou Gaudet and Drum Bay respectively.

Figure 3 and 4 give the importance values via clipped and line transect methods for Petain Lagoon. Figure 5 and 6 give the same information for Bottle Lagoon as do Figure 7 and 8 for Dead Duck Pass.

Table 1 Petain Lagoon

Density — stems/m²

| Date | 4/9 | 4/30 | 5/26 | 6/24 | 7/16 | 8/11 | 9/3 | 9/26 | 10/15 | 11/5 | 11/26 |
|------------------------|-----|------|------|------|------|------|-----|------|-------|------|-------|
| <u>Aster</u> | | | | | | | | | | | |
| <u>tenuifolius</u> | 3 | — | — | 5 | — | 6 | 6 | 1 | 5 | — | — |
| <u>Borrichia</u> | | | | | | | | | | | |
| <u>frutescens</u> | — | — | — | — | — | 8 | 6 | 13 | — | — | 1 |
| <u>Distichlis</u> | | | | | | | | | | | |
| <u>spicata</u> | 864 | 599 | 689 | 733 | 73 | 471 | 530 | 390 | 619 | 499 | 376 |
| <u>Juncus</u> | | | | | | | | | | | |
| <u>roemerianus</u> | — | — | 211 | — | — | — | — | — | — | — | — |
| <u>Salicornia</u> | | | | | | | | | | | |
| <u>virginica</u> | — | 8 | 1 | — | — | — | 29 | 1 | — | — | — |
| <u>Scirpus</u> | | | | | | | | | | | |
| <u>robustus</u> | 13 | — | 31 | 25 | 2 | 8 | 6 | — | 2 | 4 | — |
| <u>Spartina</u> | | | | | | | | | | | |
| <u>alterniflora</u> | — | 51 | — | — | 100 | — | — | — | — | — | — |
| <u>S. cynosuroides</u> | 27 | 44 | 46 | 33 | 18 | 36 | 18 | 21 | 15 | 15 | 4 |
| <u>S. patens</u> | 513 | 193 | 126 | 254 | 552 | 197 | 401 | 219 | 150 | 191 | 369 |
| | | | | | | (1) | | | | | (2) |

Also present:(1) Baccharis halimifolia(2) Fimbristylis castanea

Table 2 Petain Lagoon

Relative Density

| Date | 4/9 | 4/30 | 4/30 | 5/26 | 5/26 | 6/24 | 6/24 | 7/16 | 7/16 | 8/11 | 8/11 | 9/3 | 9/3 | 9/26 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Method | C | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT | C |
| <u>Aster</u> <u>tenuifolius</u> | .4 | - | .4 | - | - | .4 | .9 | - | - | .8 | .5 | .6 | .7 | .1 |
| <u>Borrichia</u> <u>frutescens</u> | - | - | - | - | - | - | .2 | - | - | 1.1 | .2 | .6 | .6 | 2.0 |
| <u>Distichlis</u> <u>spicata</u> | 60.8 | 67.0 | 63.4 | 62.4 | 88.1 | 69.8 | 79.0 | 9.8 | 48.9 | 64.8 | 77.1 | 53.3 | 73.4 | 60.5 |
| <u>Fimbristylis</u> <u>castanea</u> | - | - | - | - | - | - | .3 | - | - | - | - | - | .4 | - |
| <u>Juncus</u> <u>roemerianus</u> | - | - | - | 19.1 | 2.9 | - | - | - | - | - | - | - | - | - |
| <u>Salicornia</u> <u>virginica</u> | - | .8 | .7 | - | - | - | - | - | - | - | 1.5 | 2.9 | .4 | .1 |
| <u>Scirpus</u> <u>robustus</u> | .9 | - | 5.8 | 2.8 | 1.4 | 2.4 | 2.0 | .3 | 1.0 | 1.1 | .9 | .6 | .7 | - |
| <u>Spartina</u> <u>alterniflora</u> | - | 5.7 | - | - | - | - | - | 13.4 | 3.4 | - | - | - | - | - |
| <u>Spartina</u> <u>cynosuroides</u> | 1.9 | 4.9 | 26.1 | 4.3 | 3.5 | 3.1 | .5 | 2.4 | .4 | 4.9 | 1.4 | 1.8 | 1.2 | 3.3 |
| <u>Spartina</u> <u>patens</u> | 36.0 | 21.6 | 3.6 | 11.4 | 4.1 | 24.2 | 17.1 | 74.1 | 46.3 | 27.1 | 18.1 | 40.2 | 22.6 | 34.0 |

Table 2 Petain Lagoon

Relative Density

| 6/24 | 7/16 | 7/16 | 8/11 | 8/11 | 9/3 | 9/3 | 9/26 | 9/26 | 10/15 | 10/15 | 11/5 | 11/5 | 11/26 | 11/26 |
|------|------|------|------|------|------|------|------|------|-------|-------|------|------|-------|-------|
| LT | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT |
| .9 | - | - | .8 | .5 | .6 | .7 | .1 | .5 | 6.0 | .6 | - | .6 | - | - |
| .2 | - | - | 1.1 | .2 | .6 | .6 | 2.0 | 1.3 | - | - | - | - | .1 | .1 |
| 79.0 | 9.8 | 48.9 | 64.8 | 77.1 | 53.3 | 73.4 | 60.5 | 68.0 | 78.2 | 61.9 | 70.4 | 52.9 | 47.9 | 81.2 |
| .3 | - | - | - | - | - | .4 | - | - | - | 3.7 | - | 5.2 | 4.5 | - |
| - | - | - | - | - | - | - | - | - | - | 1.0 | - | - | - | - |
| - | - | - | - | 1.5 | 2.9 | .4 | .1 | 1.0 | - | .4 | - | - | - | .3 |
| 2.0 | .3 | 1.0 | 1.1 | .9 | .6 | .7 | - | .7 | 3.0 | .3 | .5 | .3 | - | - |
| - | 13.4 | 3.4 | - | - | - | - | - | - | - | - | - | - | - | - |
| .5 | 2.4 | .4 | 4.9 | 1.4 | 1.8 | 1.2 | 3.3 | .6 | 1.9 | .9 | 2.1 | .8 | .5 | .2 |
| 17.1 | 74.1 | 46.3 | 27.1 | 18.1 | 40.2 | 22.6 | 34.0 | 27.9 | 19.0 | 31.2 | 27.0 | 40.2 | 47.0 | 18.2 |

Table 3 Petain Lagoon

Relative Frequency

| Date | 4/9 | 4/30 | 4/30 | 5/26 | 5/26 | 6/24 | 6/24 | 7/16 | 7/16 | 8/11 | 8/11 | 9/3 | 9/3 | 9/3 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Method | C | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT | C |
| <u>Aster</u> <u>tenuifolius</u> | 16.7 | - | 1.7 | - | - | 6.3 | 4.2 | - | - | 11.8 | 3.1 | 9.5 | 4.7 | 5.4 |
| <u>Borrchia</u> <u>frutescens</u> | - | - | - | - | - | - | 2.1 | - | - | 5.9 | 3.1 | 9.5 | 5.4 | 11.8 |
| <u>Distichlis</u> <u>spicata</u> | 27.8 | 26.6 | 42.1 | 26.7 | 46.9 | 25.0 | 43.1 | 29.4 | 34.6 | 23.5 | 38.5 | 23.8 | 35.6 | 27.8 |
| <u>Fimbristylis</u> <u>castanea</u> | - | - | - | - | - | - | 1.2 | - | - | - | - | - | .8 | - |
| <u>Juncus</u> <u>roemerianus</u> | - | - | - | 6.7 | 3.1 | - | - | - | - | - | - | - | - | - |
| <u>Salicornia</u> <u>virginica</u> | - | 6.9 | 3.4 | 6.7 | - | - | - | - | - | - | 3.7 | 4.8 | 1.8 | 5.4 |
| <u>Scirpus</u> <u>robustus</u> | 16.7 | - | 18.0 | 26.7 | 18.8 | 25.0 | 20.8 | 11.8 | 9.8 | 11.8 | 12.8 | 14.3 | 8.9 | - |
| <u>Spartina</u> <u>alterniflora</u> | - | 26.6 | - | - | - | - | - | 23.5 | 20.3 | - | - | - | - | - |
| <u>Spartina</u> <u>cynosuroides</u> | 11.1 | 13.3 | 30.5 | 13.3 | 20.0 | 25.0 | 8.4 | 5.9 | 3.3 | 17.7 | 17.1 | 14.3 | 14.9 | 22.9 |
| <u>Spartina</u> <u>patens</u> | 27.8 | 26.6 | 4.3 | 20.0 | 11.2 | 18.8 | 20.2 | 29.4 | 32.0 | 23.5 | 21.4 | 23.8 | 27.9 | 27.8 |

Table 3 Petain Lagoon

Relative Frequency

| | 6/24 | 7/16 | 7/16 | 8/11 | 8/11 | 9/3 | 9/3 | 9/26 | 9/26 | 10/15 | 10/15 | 11/5 | 11/5 | 11/26 | 11/26 |
|---|------|------|------|------|------|------|------|------|------|-------|-------|------|------|-------|-------|
| | LT | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT |
| 4 | 4.2 | - | - | 11.8 | 3.1 | 9.5 | 4.7 | 5.5 | 2.1 | 10.1 | 3.8 | - | 3.5 | - | - |
| 3 | 2.1 | - | - | 5.9 | 3.1 | 9.5 | 5.4 | 11.1 | 7.9 | - | - | - | - | 9.1 | 2.3 |
| 0 | 43.1 | 29.4 | 34.6 | 23.5 | 38.5 | 23.8 | 35.6 | 27.8 | 40.4 | 27.8 | 45.0 | 33.3 | 41.7 | 36.3 | 68.2 |
| | 1.2 | - | - | - | - | - | .8 | - | - | - | 2.3 | - | 2.8 | 9.1 | - |
| | - | - | - | - | - | - | - | - | - | - | 1.5 | - | - | - | - |
| | - | - | - | - | 3.7 | 4.8 | 1.8 | 5.5 | 2.6 | - | .8 | - | - | - | 1.1 |
| 0 | 20.8 | 11.8 | 9.8 | 11.8 | 12.8 | 14.3 | 8.9 | - | 6.6 | 5.5 | 3.1 | 6.7 | 3.5 | - | - |
| | - | 23.5 | 20.3 | - | - | - | - | - | - | - | - | - | - | - | - |
| 0 | 8.4 | 5.9 | 3.3 | 17.7 | 17.1 | 14.3 | 14.9 | 22.2 | 10.6 | 27.8 | 12.2 | 26.7 | 12.5 | 18.2 | 3.4 |
| 8 | 20.2 | 29.4 | 32.0 | 23.5 | 21.4 | 23.8 | 27.9 | 27.8 | 29.8 | 27.8 | 31.3 | 33.3 | 36.0 | 27.3 | 25.0 |

Table 4 Petain Lagoon

Relative Dominance

| Date | 4/30 | 5/26 | 6/24 | 7/16 | 8/11 | 9/3 | 9/26 | 10/15 | 11/5 | 11/26 |
|--|------|------|------|------|------|------|------|-------|------|-------|
| <u>Aster</u> <u>tenuifolius</u> | .2 | — | .7 | — | .4 | .7 | .5 | .7 | .5 | — |
| <u>Borrichia</u> <u>frutescens</u> | — | — | .4 | — | 1.4 | 1.8 | 6.3 | — | — | .7 |
| <u>Distichlis</u> <u>spicata</u> | 51.8 | 70.3 | 72.7 | 31.0 | 67.2 | 61.0 | 51.1 | 57.9 | 53.3 | 81.4 |
| <u>Fimbristylis</u> <u>castanea</u> | — | — | — | — | — | .4 | — | 1.9 | 2.4 | — |
| <u>Juncus</u> <u>roemerianus</u> | — | 4.1 | — | — | — | — | — | 2.0 | — | — |
| <u>Salicornia</u> <u>virginica</u> | 1.2 | — | — | — | 1.3 | .4 | .8 | .3 | — | .3 |
| <u>Scirpus</u> <u>robustus</u> | 10.5 | 5.3 | 8.5 | 3.4 | 3.8 | 2.4 | 2.1 | 1.0 | 1.6 | — |
| <u>Spartina</u> <u>alterniflora</u> | — | — | — | 30.0 | — | — | — | — | — | — |
| <u>S. cynosuroides</u> | 31.2 | 18.8 | 3.4 | 3.2 | 13.1 | 10.7 | 6.3 | 9.9 | 9.0 | 2.4 |
| <u>S. patens</u> | 5.1 | 1.5 | 14.3 | 32.4 | 12.7 | 22.6 | 32.9 | 26.3 | 33.2 | 15.2 |

Table 5 Petain Lagoon

Importance Values - Clipped

| Date | 4/9 | 4/30 | 5/26 | 6/24 | 7/16 | 8/11 | 9/3 | 9/26 | 10/15 | 11/5 | 11/26 |
|--|------|------|------|------|-------|------|------|------|-------|-------|-------|
| <u>Aster</u> <u>tenuifolius</u> | 17.1 | — | — | 10.3 | — | 12.6 | 10.1 | 3.1 | 16.1 | — | — |
| <u>Borrichia</u> <u>frutescens</u> | — | — | — | — | — | 7.0 | 10.1 | 13.1 | — | — | 9.2 |
| <u>Distichlis</u> <u>spicata</u> | 88.6 | 93.6 | 89.1 | 94.8 | 39.2 | 88.3 | 77.1 | 88.3 | 106.0 | 103.4 | 84.2 |
| <u>Fimbristylis</u> <u>castanea</u> | — | — | — | — | — | — | — | — | — | — | 13.6 |
| <u>Juncus</u> <u>roemerianus</u> | — | — | 25.8 | — | — | — | — | — | — | — | — |
| <u>Salicornia</u> <u>virginica</u> | — | 7.7 | 6.7 | — | — | — | 7.7 | 5.6 | — | — | — |
| <u>Scirpus</u> <u>robustus</u> | 17.6 | — | 29.5 | 27.4 | 12.1 | 12.9 | 14.9 | — | 8.0 | 7.2 | — |
| <u>Spartina</u> <u>alterniflora</u> | — | 32.3 | — | — | 36.9 | — | — | — | — | — | — |
| <u>S. cynosuroides</u> | 13.0 | 18.2 | 17.6 | 28.1 | 8.3 | 22.6 | 16.1 | 22.5 | 29.7 | 28.8 | 18.7 |
| <u>S. patens</u> | 63.8 | 58.2 | 31.4 | 42.9 | 103.5 | 50.6 | 64.0 | 61.8 | 46.8 | 60.3 | 74.3 |

Table 6 Petain Lagoon

Importance Values — Line Transect

| Date | 4/30 | 5/26 | 6/24 | 7/16 | 8/11 | 9/3 | 9/26 | 10/15 | 11/5 | 11/26 |
|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <u>Aster</u> <u>tenuifolius</u> | 2.3 | — | 5.8 | — | 4.0 | 6.1 | 3.1 | 5.1 | 4.6 | — |
| <u>Borrichia</u> <u>frutescens</u> | — | — | 2.7 | — | 4.7 | 7.8 | 15.5 | — | — | 3.1 |
| <u>Distichlis</u> <u>spicata</u> | 157.3 | 205.3 | 194.8 | 114.5 | 182.8 | 170.0 | 159.5 | 164.8 | 147.9 | 230.8 |
| <u>Fimbristylis</u> <u>castanea</u> | — | — | 1.5 | — | — | 1.6 | — | 7.9 | 10.4 | — |
| <u>Juncus</u> <u>roemerianus</u> | — | 10.1 | — | — | — | — | — | 4.5 | — | — |
| <u>Salicornia</u> <u>virginica</u> | 5.3 | — | — | — | 6.5 | 2.6 | 4.8 | 1.5 | — | 1.7 |
| <u>Scirpus</u> <u>robustus</u> | 34.3 | 25.5 | 31.3 | 14.2 | 17.5 | 12.0 | 9.4 | 4.4 | 5.4 | — |
| <u>Spartina</u> <u>alterniflora</u> | — | — | — | 53.7 | — | — | — | — | — | — |
| <u>S. cynosuroides</u> | 87.8 | 42.3 | 12.3 | 6.9 | 31.6 | 16.9 | 17.5 | 22.8 | 22.3 | 6.0 |
| <u>S. patens</u> | 13.0 | 16.8 | 51.6 | 110.7 | 52.2 | 73.1 | 90.6 | 88.8 | 109.4 | 58.2 |

Table 7 Bayou Gaudet

Density - Stems/m²

| Date | 4/14 | 5/13 | 6/13 | 7/1 | 7/30 | 9/3 | 9/26 | 10/15 | 11/5 | 11/26 | 1/15 |
|-------------------------------------|------|------|------|------|------|------|------|-------|------|-------|------|
| <u>Distichlis</u> <u>spicata</u> | 44 | 258 | 48 | 21 | 20 | 21 | 21 | 32 | 16 | 19 | - |
| <u>Spartina</u> <u>patens</u> | 1511 | 1280 | 1980 | 1610 | 1495 | 1775 | 1308 | 1193 | 1340 | 1228 | 1438 |

Table 8 Bayou Gaudet

Relative Density

| Date | 4/14 | 4/16 | 5/13 | 5/13 | 5/26 | 6/13 | 6/13 | 7/1 | 7/30 | 9/3 | 9/26 | 10/15 | 11/5 | 11/26 | 1/15 |
|-------------------------------------|------|------|------|------|-----------|------|-----------|------|------|------|------|-------|------|-------|------|
| Method | C | LT | C | LT | LT | C | LT | C | C | C | C | C | C | C | C |
| <u>Distichlis</u> <u>spicata</u> | 2.9 | - | 16.7 | 23.6 | .6 | 2.5 | 10.6 | 1.2 | 1.3 | 1.1 | 1.5 | 2.5 | 1.1 | 1.6 | - |
| <u>Spartina</u> <u>patens</u> | 97.1 | 100 | 83.3 | 76.4 | 99.4 * | 97.5 | 89.4 * | 98.8 | 98.7 | 98.9 | 98.5 | 97.5 | 98.9 | 98.4 | 100 |

Also present:*Scirpus olneyi

Table 9 Bayou Gaudet

Relative Frequency

| Date | 4/14 | 4/16 | 5/13 | 5/13 | 5/26 | 6/13 | 6/13 | 7/1 | 7/30 | 9/3 | 9/26 | 10/15 | 11/5 | 11/26 | 1/15 |
|-------------------------------------|------|------|------|------|------|------|------|------|------|------|------|-------|------|-------|------|
| Method | C | LT | C | LT | LT | C | LT | C | C | C | C | C | C | C | C |
| <u>Distichlis</u> <u>spicata</u> | 42.9 | - | 50.0 | 45.5 | 9.9 | 50.0 | 24.5 | 25.0 | 40.0 | 50.0 | 39.8 | 40.0 | 40.0 | 25.0 | - |
| <u>Spartina</u> <u>patens</u> | 57.1 | 100 | 50.0 | 54.5 | 88.4 | 50.0 | 73.4 | 75.0 | 60.0 | 50.0 | 60.2 | 60.0 | 60.0 | 75.0 | 100 |

Table 10 Bayou Gaudet

Relative Dominance

| Date | 4/16 | 5/13 | 5/20 | 6/13 |
|-------------------------------------|------|------|------|------|
| <u>Distichlis</u> <u>spicata</u> | - | 32.0 | 1.1 | 11.3 |
| <u>Spartina</u> <u>patens</u> | 100 | 68.0 | 98.5 | 85.9 |

Table 11 Bayou Gaudet

Importance Values - Clipped

| Date | 4/14 | 5/13 | 6/13 | 7/1 | 7/30 | 9/3 | 9/26 | 10/15 | 11/5 | 11/26 | 1/15 |
|-------------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|
| <u>Distichlis</u> <u>spicata</u> | 45.8 | 66.7 | 52.5 | 26.2 | 41.3 | 51.1 | 41.3 | 42.5 | 41.1 | 26.6 | - |
| <u>Spartina</u> <u>patens</u> | 154.2 | 133.3 | 147.5 | 173.8 | 158.7 | 148.9 | 158.7 | 157.5 | 158.9 | 173.4 | 200 |

Table 12 Bayou Gaudet

Importance Values - Line Transect

| Date | 4/16 | 5/13 | 5/26 | 6/13 |
|-------------------------------------|------|-------|-------|-------|
| <u>Distichlis</u> <u>spicata</u> | - | 101.1 | 11.6 | 46.4 |
| <u>Spartina</u> <u>patens</u> | 300 | 198.9 | 286.3 | 248.7 |

Table 13 Bottle Lagoon

Density — stems/m²

| Date | 6/10 | 6/27 | 7/22 | 8/11 | 9/10 | 10/3 | 10/23 | 11/12 | 12/2 |
|--|------|------|------|------|------|------|-------|-------|------|
| <u>Aster</u> <u>tenuifolius</u> | 13 | 15 | 14 | 8 | 1 | 4 | 7 | 6 | — |
| <u>Borrichia</u> <u>frutescens</u> | 5 | — | — | — | — | 2 | — | — | — |
| <u>Distichlis</u> <u>spicata</u> | 694 | 605 | 525 | 404 | 807 | 498 | 135 | 441 | 402 |
| <u>Juncus</u> <u>roemerianus</u> | 79 | 30 | 59 | 159 | 92 | 162 | 202 | 146 | 170 |
| <u>Salicornia</u> <u>virginica</u> | 18 | — | 2 | — | 6 | 55 | — | 19 | — |
| <u>Scirpus</u> <u>robustus</u> | 5 | 7 | 14 | 3 | — | — | — | — | — |
| <u>Spartina</u> <u>alterniflora</u> | 87 | 115 | 138 | 155 | 83 | 100 | 151 | 113 | 54 |
| <u>S. patens</u> | 118 | 290 | 107 | 76 | 134 | 182 | 98 | 340 | 231 |
| | | | (1) | (2) | | | | (3) | |

Also present:

(1) Agalinis maritima(2) Eleocharis sp.(3) Lythrum lineare

Table 14 Bottle Lagoon

Relative Density

| Date | 6/10 | 6/10 | 6/27 | 6/27 | 7/22 | 7/22 | 8/11 | 8/11 | 9/10 | 9/10 | 10/3 | 10/ |
|--|------|------|------|------|------|------|------|------|------|------|------|-----|
| Method | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT |
| <u>Aster</u> <u>tenuifolius</u> | 1.2 | .5 | 1.4 | .7 | 1.6 | 1.1 | 1.0 | .6 | - | .1 | 1.0 | . |
| <u>Borrichia</u> <u>frutescens</u> | .5 | .4 | - | 1.0 | - | .2 | - | - | - | - | .3 | - |
| <u>Distichlis</u> <u>spicata</u> | 68.1 | 78.4 | 56.9 | 60.0 | 61.0 | 74.4 | 50.2 | 48.6 | 72.0 | 63.2 | 49.8 | 61. |
| <u>Juncus</u> <u>roemerianus</u> | 7.8 | 9.1 | 2.8 | 7.3 | 6.8 | 7.3 | 19.8 | 20.6 | 8.2 | 14.2 | 16.2 | 9. |
| <u>Salicornia</u> <u>virginica</u> | 1.8 | 1.1 | - | .3 | .2 | 2.2 | - | - | .4 | .1 | 5.5 | - |
| <u>Scirpus</u> <u>robustus</u> | .5 | .2 | .8 | .2 | 1.6 | .4 | .4 | .6 | - | .1 | - | - |
| <u>Spartina</u> <u>alterniflora</u> | 8.5 | 4.7 | 10.8 | 4.2 | 16.2 | 6.6 | 19.2 | 14.7 | 7.4 | 6.5 | 10.0 | 7. |
| <u>Spartina</u> <u>patens</u> | 11.6 | 5.6 | 27.3 | 27.1 | 12.4 | 7.8 | 9.4 | 14.9 | 12.0 | 15.9 | 18.2 | 20. |

FOLLOUT FRAME /

Table 14 Bottle Lagoon

Relative Density

| 7/22 | 7/22 | 8/11 | 8/11 | 9/10 | 9/10 | 10/3 | 10/3 | 10/23 | 10/23 | 11/12 | 11/12 | 12/2 | 12/2 |
|------|------|------|------|------|------|------|------|-------|-------|-------|-------|------|------|
| C | LT | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT |
| 1.6 | 1.1 | 1.0 | .6 | - | .1 | 1.0 | .5 | 1.1 | - | .6 | 1.6 | - | .5 |
| - | .2 | - | - | - | - | .3 | - | - | - | - | .1 | - | .5 |
| 61.0 | 74.4 | 50.2 | 48.6 | 72.0 | 63.2 | 49.8 | 61.3 | 22.8 | 50.0 | 41.4 | 55.1 | 46.9 | 52.2 |
| 6.8 | 7.3 | 19.8 | 20.6 | 8.2 | 14.2 | 16.2 | 9.6 | 34.1 | 15.1 | 13.7 | 15.0 | 19.8 | 14.2 |
| .2 | 2.2 | - | - | .4 | .1 | 5.5 | - | - | 1.8 | 1.8 | 1.1 | - | - |
| 1.6 | .4 | .4 | .6 | - | .1 | - | - | - | .5 | - | - | - | - |
| 16.2 | 6.6 | 19.2 | 14.7 | 7.4 | 6.5 | 10.0 | 7.7 | 25.5 | 19.5 | 10.6 | 15.1 | 6.3 | 12.0 |
| 12.4 | 7.8 | 9.4 | 14.9 | 12.0 | 15.9 | 18.2 | 20.9 | 16.5 | 13.1 | 31.9 | 9.3 | 27.0 | 20.6 |

Table 15 Bottle Lagoon

Relative Frequency

| Date | 6/10 | 6/10 | 6/27 | 6/27 | 7/22 | 7/22 | 8/11 | 8/11 | 9/10 | 9/10 | 10/3 | 10/3 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|
| Method | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT |
| <u>Aster</u> <u>tenuifolius</u> | 8.3 | 1.9 | 10.0 | 3.4 | 10.0 | 2.2 | 6.6 | 2.1 | 5.9 | 1.0 | 10.5 | 2.2 |
| <u>Borrichia</u> <u>frutescens</u> | 4.3 | 1.9 | - | - | - | .8 | - | - | - | - | 5.3 | - |
| <u>Distichlis</u> <u>spicata</u> | 25.0 | 40.6 | 30.0 | 36.5 | 25.0 | 38.4 | 26.7 | 25.0 | 29.4 | 36.8 | 26.3 | 34.5 |
| <u>Juncus</u> <u>roemerianus</u> | 4.3 | 8.5 | 5.1 | 6.9 | 5.0 | 8.4 | 13.4 | 11.4 | 5.9 | 9.4 | 5.3 | 8.8 |
| <u>Salicornia</u> <u>virginica</u> | 8.3 | 2.2 | - | 1.7 | 5.0 | 3.9 | - | - | 11.8 | 2.0 | 5.3 | - |
| <u>Scirpus</u> <u>robustus</u> | 8.3 | 1.9 | 10.0 | 2.2 | 10.0 | 2.2 | 6.6 | 2.1 | - | 1.0 | - | - |
| <u>Spartina</u> <u>alterniflora</u> | 20.8 | 32.1 | 24.9 | 29.7 | 25.0 | 32.4 | 26.7 | 43.5 | 23.5 | 34.1 | 21.0 | 37.7 |
| <u>Spartina</u> <u>patens</u> | 20.8 | 10.9 | 20.0 | 19.6 | 15.0 | 11.7 | 20.0 | 14.3 | 23.5 | 14.7 | 26.3 | 16.8 |

Table 15 Bottle Lagoon

Relative Frequency

| | 7/22 | 8/11 | 8/11 | 9/10 | 9/10 | 10/3 | 10/3 | 10/23 | 10/23 | 11/12 | 11/12 | 12/2 | 12/2 |
|-----|------|------|------|------|------|------|------|-------|-------|-------|-------|------|------|
| | LT | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT |
| 0.0 | 2.2 | 6.6 | 2.1 | 5.9 | 1.0 | 10.5 | 2.2 | 6.3 | - | 5.6 | 4.6 | - | 1.0 |
| | .8 | - | - | - | - | 5.3 | - | - | - | - | .9 | - | 1.0 |
| 5.0 | 38.4 | 26.7 | 25.0 | 29.4 | 36.8 | 26.3 | 34.5 | 31.2 | 29.6 | 27.8 | 32.8 | 33.3 | 37.8 |
| 5.0 | 8.4 | 13.4 | 11.4 | 5.9 | 9.4 | 5.3 | 8.8 | 6.3 | 8.7 | 5.6 | 9.9 | 11.1 | 7.6 |
| 5.0 | 3.9 | - | - | 11.8 | 2.0 | 5.3 | - | - | .8 | 11.1 | 1.5 | - | - |
| 5.0 | 2.2 | 6.6 | 2.1 | - | 1.0 | - | - | - | .8 | - | - | - | - |
| 5.0 | 32.4 | 26.7 | 43.5 | 23.5 | 34.1 | 21.0 | 37.7 | 31.2 | 50.5 | 22.2 | 39.6 | 22.3 | 37.0 |
| 5.0 | 11.7 | 20.0 | 14.3 | 23.5 | 14.7 | 26.3 | 16.8 | 25.0 | 9.6 | 27.8 | 8.4 | 33.3 | 15.1 |

FOLDOUT FRAME 2

Table 16 Bottle Lagoon

Relative Dominance

| Date | 6/10 | 6/27 | 7/22 | 8/11 | 9/10 | 10/3 | 10/23 | 11/12 | 12/2 |
|--|------|------|------|------|------|------|-------|-------|------|
| <u>Aster</u> <u>tenuifolius</u> | .3 | 1.2 | .5 | .2 | .2 | .6 | — | 1.0 | .4 |
| <u>Borrchia</u> <u>frutescens</u> | .4 | .1 | — | — | — | — | — | .3 | 1.5 |
| <u>Distichlis</u> <u>spicata</u> | 49.4 | 39.8 | 36.5 | 15.8 | 33.6 | 29.6 | 13.7 | 19.7 | 21.1 |
| <u>Juncus</u> <u>roemerianus</u> | 11.7 | 9.9 | 6.1 | 11.6 | 11.8 | 7.2 | 6.6 | 12.9 | 13.7 |
| <u>Salicornia</u> <u>virginica</u> | .8 | .4 | 1.1 | — | .3 | — | .5 | .2 | — |
| <u>Scirpus</u> <u>robustus</u> | .5 | .6 | .7 | .8 | .3 | — | .2 | — | — |
| <u>Spartina</u> <u>alterniflora</u> | 33.9 | 31.6 | 52.6 | 67.5 | 47.0 | 52.5 | 75.4 | 62.1 | 55.8 |
| <u>S. patens</u> | 3.1 | 16.4 | 2.7 | 4.0 | 6.8 | 10.1 | 3.6 | 3.0 | 7.5 |

Table 17 Bottle Lagoon

| <u>Importance Values — Clipped</u> | | | | | | | | | |
|--|------|------|------|------|-------|------|-------|-------|------|
| Date | 6/10 | 6/27 | 7/22 | 8/11 | 9/10 | 10/3 | 10/23 | 11/12 | 12/2 |
| <u>Aster</u> <u>tenuifolius</u> | 9.5 | 11.4 | 11.6 | 7.6 | 5.9 | 11.5 | 7.4 | 6.2 | — |
| <u>Borrichia</u> <u>frutescens</u> | 4.8 | — | — | — | — | 5.6 | — | — | — |
| <u>Distichlis</u> <u>spicata</u> | 93.1 | 86.9 | 86.0 | 76.9 | 101.4 | 76.1 | 54.0 | 69.2 | 80.2 |
| <u>Juncus</u> <u>roemerianus</u> | 12.1 | 7.9 | 11.8 | 33.2 | 14.1 | 21.5 | 40.4 | 19.3 | 30.9 |
| <u>Salicornia</u> <u>virginica</u> | 10.1 | — | 5.2 | — | 12.2 | 10.8 | — | 12.9 | — |
| <u>Scirpus</u> <u>robustus</u> | 8.8 | 10.8 | 11.6 | 7.0 | — | — | — | — | — |
| <u>Spartina</u> <u>alterniflora</u> | 29.3 | 35.7 | 41.2 | 45.9 | 30.9 | 31.0 | 56.7 | 32.8 | 28.6 |
| <u>S. patens</u> | 32.4 | 47.3 | 27.4 | 29.4 | 35.5 | 44.5 | 41.5 | 59.7 | 60.3 |

Table 18 Bottle Lagoon

Importance Values — Line Transect

| Date | 6/10 | 6/27 | 7/22 | 8/11 | 9/10 | 10/3 | 10/23 | 11/12 | 12/2 |
|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <u>Aster</u> <u>tenuifolius</u> | 2.7 | 5.3 | 3.8 | 2.9 | 1.3 | 3.3 | — | 7.0 | 1.9 |
| <u>Borrichia</u> <u>frutescens</u> | 2.7 | .2 | 1.0 | — | — | — | — | 1.3 | 3.0 |
| <u>Distichlis</u> <u>spicata</u> | 168.4 | 136.3 | 149.5 | 89.4 | 133.6 | 125.4 | 93.3 | 107.6 | 111.1 |
| <u>Juncus</u> <u>roemerianus</u> | 29.3 | 24.1 | 21.8 | 43.6 | 34.8 | 25.6 | 30.4 | 37.8 | 35.5 |
| <u>Salicornia</u> <u>virginica</u> | 4.1 | 2.4 | 7.2 | — | 2.4 | — | 3.1 | 2.8 | — |
| <u>Scirpus</u> <u>robustus</u> | 2.6 | 3.0 | 3.3 | 3.5 | 1.4 | — | 1.5 | — | — |
| <u>Spartina</u> <u>alterniflora</u> | 70.7 | 65.5 | 91.6 | 125.7 | 87.6 | 97.9 | 145.4 | 116.8 | 104.8 |
| <u>S. patens</u> | 19.6 | 63.1 | 22.2 | 33.2 | 37.4 | 47.8 | 26.3 | 20.7 | 43.2 |

Table 19 Dead Duck Pass

| Date | <u>Density — stems/m²</u> | | | | | | | | | | | | |
|--|--------------------------------------|------|------|------|-----|------|------|-----|--------|-------|-------|------|-----|
| | 3/27 | 4/23 | 5/21 | 6/16 | 7/9 | 7/29 | 8/27 | 9/8 | 10/3 | 10/23 | 11/12 | 12/2 | 2/5 |
| <u>Aster</u> <u>tenuifolius</u> | — | 13 | 10 | 6 | — | 14 | 1 | 26 | 2 | 3 | — | 4 | 1 |
| <u>Borrichia</u> <u>frutescens</u> | — | — | 2 | — | 13 | — | — | 4 | 4 | 3 | — | — | — |
| <u>Distichlis</u> <u>spicata</u> | 61 | 34 | 84 | 81 | 15 | 245 | 16 | 33 | 12 | 8 | 22 | 23 | 13 |
| <u>Juncus</u> <u>roemerianus</u> | 472 | 423 | 878 | 673 | 663 | 485 | 722 | 475 | 511 | 261 | 350 | 586 | 486 |
| <u>Spartina</u> <u>alterniflora</u> | 286 | 113 | 46 | 90 | 110 | 97 | 84 | 68 | 134 | 157 | 160 | 82 | 130 |
| <u>S. patens</u> | 6 | 13 | 16 | 78 | 35 | — | 88 | 182 | 8 | — | 1 | — | 1 |
| | | | (1) | (2) | (2) | | | | (1)(3) | (3) | (1) | | |

Also present:

- (1) Salicornia virginica
- (2) Eleocharis sp.
- (3) Batis maritima

Table 20 Dead Duck Pass

Relative Density

| Date | 3/27 | 4/16 | 4/23 | 4/23 | 5/21 | 5/21 | 6/16 | 6/16 | 7/9 | 7/9 | 7/29 | 7/29 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|
| Method | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT |
| <u>Aster</u> <u>tenuifolius</u> | - | 1.1 | 2.2 | 1.7 | .9 | .2 | .7 | .4 | - | 1.1 | 1.8 | 1.3 |
| <u>Borrichia</u> <u>frutescens</u> | - | 2.6 | - | .5 | .2 | .7 | - | - | 1.6 | .2 | - | - |
| <u>Distichlis</u> <u>spicata</u> | 7.4 | 25.4 | 5.7 | 4.5 | 8.1 | 6.6 | 8.7 | 4.3 | 1.8 | 6.3 | 29.1 | 20.5 |
| <u>Juncus</u> <u>roemerianus</u> | 57.2 | 63.0 | 71.0 | 59.4 | 84.8 | 83.3 | 72.5 | 74.1 | 79.3 | 78.2 | 57.6 | 64.5 |
| <u>Spartina</u> <u>alterniflora</u> | 34.6 | 7.9 | 18.9 | 12.3 | 4.4 | 8.5 | 9.7 | 10.0 | 13.1 | 10.4 | 11.5 | 11.9 |
| <u>Spartina</u> <u>patens</u> | .8 | - | 2.2 | 21.6 | 1.5 | .3 | 8.7 | 11.2 | 4.2 | 3.8 | - | 1.8 |

| 8/27 | 8/27 | 9/8 | 10/3 | 10/3 | 10/23 | 10/23 | 11/12 | 11/12 | 12/2 | 12/2 | 2/5 | 2/5 |
|------|------|------|------|------|-------|-------|-------|-------|------|------|------|------|
| C | LT | C | C | LT | C | LT | C | LT | C | LT | C | LT |
| - | - | 3.3 | .3 | - | .7 | 1.1 | - | .6 | .5 | - | .2 | - |
| - | .8 | .5 | .5 | - | .7 | - | - | - | - | - | - | .7 |
| 1.7 | 5.0 | 4.2 | 1.8 | 1.3 | 1.8 | 7.6 | 4.1 | 2.6 | 3.3 | 1.3 | 2.1 | 2.0 |
| 79.4 | 70.0 | 60.3 | 75.8 | 75.1 | 60.2 | 73.7 | 65.7 | 62.9 | 84.4 | 78.6 | 76.9 | 65.9 |
| 9.2 | 14.2 | 8.6 | 19.9 | 16.5 | 36.2 | 12.8 | 30.0 | 21.9 | 11.8 | 17.5 | 20.6 | 30.7 |
| 9.7 | 10.2 | 23.1 | 1.2 | 3.7 | - | 4.8 | .2 | 10.5 | - | 2.6 | .2 | .7 |

Table 21 Dead Duck Pass

Relative Frequency

| Date | 3/27 | 4/16 | 4/23 | 4/23 | 5/21 | 5/21 | 6/16 | 6/16 | 7/9 | 7/9 | 7/29 | 7/29 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|
| Method | C | LT | C | LT | C | LT | C | LT | C | LT | C | LT |
| <u>Aster</u> <u>tenuifolius</u> | - | 3.4 | 8.3 | 3.9 | 7.1 | 1.1 | 15.0 | 1.2 | - | 2.1 | 14.3 | 4.0 |
| <u>Borrichia</u> <u>frutescens</u> | - | 13.4 | - | 5.2 | 7.1 | 6.7 | - | - | 10.0 | 1.1 | - | - |
| <u>Distichlis</u> <u>spicata</u> | 30.0 | 20.0 | 25.0 | 9.5 | 28.6 | 13.2 | 20.0 | 7.7 | 20.0 | 7.5 | 28.6 | 16.8 |
| <u>Juncus</u> <u>roemerianus</u> | 23.3 | 33.2 | 16.7 | 25.6 | 21.5 | 39.2 | 20.0 | 39.4 | 20.0 | 38.3 | 21.3 | 34.9 |
| <u>Spartina</u> <u>alterniflora</u> | 33.3 | 30.0 | 33.3 | 34.2 | 21.5 | 35.3 | 25.0 | 45.1 | 40.0 | 46.8 | 35.8 | 43.0 |
| <u>Spartina</u> <u>patens</u> | 10.0 | - | 16.7 | 21.6 | 7.1 | 3.2 | 20.0 | 4.4 | 10.0 | 2.1 | - | 1.3 |

Table 21 (cont.)

| 8/27 | 8/27 | 9/8 | 10/3 | 10/3 | 10/23 | 10/23 | 11/12 | 11/12 | 12/2 | 12/2 | 2/5 | 2/5 |
|------|------|------|------|------|-------|-------|-------|-------|------|------|------|------|
| C | LT | C | C | LT | C | LT | C | LT | C | LT | C | LT |
| 9.3 | - | 6.7 | 7.7 | - | 8.3 | 1.3 | - | 1.3 | 8.3 | - | 7.1 | - |
| - | 1.4 | 6.7 | 7.7 | - | 8.3 | - | - | - | - | - | - | 2.3 |
| 9.1 | 4.7 | 26.6 | 15.4 | 1.0 | 25.0 | 7.0 | 20.0 | 2.5 | 25.0 | 2.5 | 28.6 | 2.3 |
| 27.3 | 43.5 | 20.0 | 23.1 | 38.8 | 16.7 | 38.8 | 20.0 | 35.0 | 25.0 | 35.9 | 28.6 | 37.2 |
| 27.3 | 45.7 | 20.0 | 30.7 | 55.8 | 33.4 | 49.4 | 50.0 | 54.9 | 41.7 | 59.0 | 28.6 | 57.0 |
| 27.3 | 4.7 | 20.0 | 7.7 | 2.2 | - | 3.5 | 10.0 | 5.0 | - | 2.5 | 7.1 | 1.2 |

Table 22 Dead Duck Pass

| Date | <u>Relative Dominance</u> | | | | | | | | | | | |
|--|---------------------------|------|------|------|------|------|------|------|-------|-------|------|------|
| | 4/16 | 4/23 | 5/21 | 6/16 | 7/9 | 7/29 | 8/27 | 10/3 | 10/23 | 11/12 | 12/2 | 2/5 |
| <u>Aster</u> <u>tenuifolius</u> | .7 | .7 | .1 | .1 | .6 | .4 | — | — | .3 | .1 | — | — |
| <u>Borrichia</u> <u>frutescens</u> | 7.3 | 1.2 | 1.8 | — | .3 | — | .3 | — | — | — | — | .4 |
| <u>Distichlis</u> <u>spicata</u> | 12.8 | 2.4 | 3.1 | 1.5 | 2.2 | 6.6 | 1.2 | .4 | 2.4 | .7 | .2 | .4 |
| <u>Juncus</u> <u>roemerianus</u> | 55.3 | 52.4 | 70.3 | 52.1 | 55.6 | 40.5 | 58.5 | 47.7 | 52.4 | 36.2 | 37.2 | 48.4 |
| <u>Spartina</u> <u>alterniflora</u> | 24.9 | 36.2 | 23.6 | 39.3 | 40.1 | 52.2 | 38.1 | 50.2 | 43.7 | 60.6 | 62.2 | 50.7 |
| <u>S. patens</u> | — | 7.1 | .9 | 3.5 | 1.2 | .4 | 1.9 | .8 | 1.2 | 2.0 | .4 | .1 |

Table 23 Dead Duck Pass

Importance Values - Clipped

| Date | 3/27 | 4/23 | 5/21 | 6/16 | 7/9 | 7/29 | 8/27 | 9/8 | 10/3 | 10/23 | 11/12 | 12/2 | 2/5 |
|--|------|------|-------|------|------|------|-------|------|------|-------|-------|-------|-------|
| <u>Aster</u> <u>tenuifolius</u> | — | 10.5 | 8.0 | 15.7 | — | 16.1 | 9.3 | 10.0 | 8.0 | 9.0 | — | 8.8 | 7.3 |
| <u>Borrichia</u> <u>frutescens</u> | — | — | 7.3 | — | 11.6 | — | — | 7.2 | 8.2 | 9.0 | — | — | — |
| <u>Distichlis</u> <u>spicata</u> | 37.4 | 30.0 | 36.7 | 28.7 | 21.8 | 57.7 | 10.8 | 30.8 | 17.2 | 26.8 | 24.1 | 28.3 | 30.7 |
| <u>Juncus</u> <u>roemerianus</u> | 80.5 | 87.7 | 106.3 | 92.5 | 99.3 | 78.9 | 106.7 | 80.3 | 98.9 | 76.9 | 85.7 | 109.4 | 105.5 |
| <u>Spartina</u> <u>alterniflora</u> | 67.9 | 52.2 | 25.9 | 34.7 | 53.1 | 47.3 | 36.5 | 28.6 | 50.6 | 69.6 | 80.0 | 53.5 | 49.2 |
| <u>S. patens</u> | 10.8 | 18.9 | 8.6 | 28.7 | 14.2 | — | 37.0 | 43.1 | 8.9 | — | 10.2 | — | 7.3 |

Table 24 Dead Duck PassImportance Values — Line Transect

| Date | 4/16 | 4/23 | 5/21 | 6/16 | 7/9 | 7/29 | 8/27 | 10/3 | 10/23 | 11/12 | 12/2 | 2/5 |
|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <u>Aster</u> <u>tenuifolius</u> | 5.2 | 6.3 | 1.3 | 1.7 | 3.8 | 5.6 | — | — | 2.7 | 2.0 | — | — |
| <u>Borrichia</u> <u>frutescens</u> | 23.3 | 6.9 | 8.8 | — | 1.6 | — | 2.5 | — | — | — | — | 3.4 |
| <u>Distichlis</u> <u>spicata</u> | 58.2 | 16.4 | 22.9 | 13.5 | 16.0 | 43.9 | 10.9 | 2.7 | 17.0 | 5.8 | 4.0 | 6.7 |
| <u>Juncus</u> <u>roemerianus</u> | 151.5 | 137.4 | 192.8 | 165.6 | 172.1 | 139.9 | 172.0 | 161.6 | 164.9 | 134.1 | 151.7 | 151.5 |
| <u>Spartina</u> <u>alterniflora</u> | 62.8 | 82.7 | 67.4 | 94.4 | 97.3 | 107.1 | 98.0 | 122.5 | 105.9 | 137.4 | 138.7 | 138.4 |
| <u>S. patens</u> | — | 50.3 | 4.4 | 19.1 | 7.1 | 3.4 | 16.8 | 6.7 | 9.5 | 17.5 | 5.5 | 2.0 |

Table 25 Drum Bay

Density — stems/m²

| Date | 3/25 | 4/16 | 5/15 | 6/19 | 7/10 | 7/30 | 8/28 | 9/17 | 10/8 | 10/29 | 11/19 | 12/2 | 12/10 | 12/15 | 1/5 |
|------------------------------|---------|------|------|------|------|------|------|------|------|--------------|-------|------|-------|-------|-----|
| <u>Spartina alterniflora</u> | 473 | 440 | 442 | 438 | 441 | 437 | 469 | 470 | 494 | (653) 592 | 423 | 621 | 520 | 267 | 525 |
| | (1) (2) | | | | | | | (1) | | (2) | (a) | | | | |

Also present:

- (1) Distichlis spicata
- (2) Avicennia nitida seedlings

(a) This sample is not consistent with others due to inexperienced sorter.

Figure 1. Density of Spartina patens in stems/m² at Bayou Gaudet site versus week of the year 1975.

DENSITY




Figure 2. Density of Spartina alterniflora in stems/m²
at Drum Bay site versus week of the years
1975 and early 1976.

DENSITY









Figure 3. Clipped importance values from Petain Lagoon site versus week of the year 1975.

| | |
|-------------------------------------|---|
| <u>Distichlis</u> <u>spicata</u> |  |
| <u>Scirpus</u> <u>robustus</u> |  |
| <u>Spartina</u> <u>cynosuroides</u> |  |
| <u>Spartina</u> <u>patens</u> |  |











Figure 4. Line transect importance values from Petain Lagoon site versus week of the year 1975.

| | |
|------------------------------|---|
| <u>Distichlis spicata</u> |  |
| <u>Scirpus robustus</u> |  |
| <u>Spartina cynosuroides</u> |  |
| <u>Spartina patens</u> |  |

IMP.
VAL.










Figure 5. Clipped importance values from Bottle Lagoon
versus week of the year 1975.

| | |
|------------------------------|---|
| <u>Distichlis spicata</u> |  |
| <u>Juncus roemerianus</u> |  |
| <u>Spartina alterniflora</u> |  |
| <u>Spartina patens</u> |  |








Figure 6. Line transect importance values from Bottle Lagoon versus week of the year 1975.

| | |
|------------------------------|---|
| <u>Distichlis spicata</u> |  |
| <u>Juncus roemerianus</u> |  |
| <u>Spartina alterniflora</u> |  |
| <u>Spartina patens</u> |  |






Figure 7. Clipped importance values from Dead Duck Pass versus week of the years 1975 and early 1976.














Figure 8. Line transect importance values from Dead Duck Pass versus week of the years 1975 and early 1976.

| | |
|------------------------------|--|
| <u>Distichlis spicata</u> |  |
| <u>Juncus roemerianus</u> |  |
| <u>Spartina alterniflora</u> |  |
| <u>Spartina patens</u> |  |





WEEK

DISCUSSION

Introduction

This study involved an analysis of marsh vegetation over a one year period to determine seasonal variation in the species composition and its relationship to salinity. Two methods of sampling the vegetation were utilized, the clipped method and the line transect. This was done to more adequately survey the vegetation and to make a comparison of methods as applied to marsh vegetation. In addition, salinity regimes were established and vegetation surveyed in each regime.

Salinity

From the values obtained as presented in Table 26 salinity regimes can be established based on peak soil salinities (PSS). Petain Lagoon and Bayou Gaudet exhibit PSS of 10 ppt or less, Bottle Lagoon between 10 and 20 ppt, Dead Duck Pass between 15 and 25 ppt and Drum Bay above 20 ppt with regular flooding.

In correlating salinity with vegetation, Spartina patens occurs as the dominant species or in pure stands when the PSS is 10 ppt or less. The vegetation of Petain Lagoon is in a successional sequence toward pure S. patens. This vegetation is also found in areas with a PSS of 10 ppt or less.

In those areas such as Bottle Lagoon in which the PSS is between 10 and 20 ppt, Distichlis spicata is the dominant species throughout

Table 26 Salinity - ‰

Petain Lagoon

| Date | 4/9 | 5/26 | 6/24 | 8/11 | 9/13 | 9/26 | 10/15 | 11/5 | 10/26 |
|------------|------|---------|---------|---------|---------|------|---------|---------|---------|
| Water type | soil | surface | surface | surface | surface | soil | surface | surface | surface |
| Salinity | 9.0 | 3.5 | 2.0 | 7.0 | 5.0 | 7.5 | 4.0 | 4.0 | 0.1 |

Bayou Gaudet

| Date | 4/14 | 4/16 | 5/13 | 5/21 | 6/13 | 7/1 | 9/3 | 9/26 | 10/15 | 11/5 |
|------------|---------|------|---------|------|------|---------|---------|---------|---------|---------|
| Water type | surface | soil | surface | soil | soil | surface | surface | surface | surface | surface |
| Salinity | 6.0 | 8.5 | 5.0 | 9.0 | 8.5 | 5.0 | 5.5 | 5.0 | 4.0 | 5.0 |

Bottle Lagoon

| Date | 6/10 | 6/27 | 7/1 | 7/22 | 8/11 | 9/10 | 10/3 | 10/23 | 11/12 |
|------------|---------|------|------|------|---------|---------|---------|---------|---------|
| Water type | surface | soil | soil | soil | surface | surface | surface | surface | surface |
| Salinity | 3.75 | 7.0 | 14.0 | 19.0 | 8.75 | 5.0 | 5.5 | 3.5 | 3.0 |

Dead Duck Pass

| Date | 4/16 | 4/23 | 5/21 | 6/16 | 7/9 | 7/29 | 8/27 | 10/3 | 10/23 | 11/12 |
|------------|------|------|------|---------|------|---------|---------|---------|-------|---------|
| Water type | soil | soil | soil | surface | soil | surface | surface | surface | soil | surface |
| Salinity | 10.0 | 8.5 | 19.0 | 5.0 | 23.5 | 7.25 | 5.5 | 5.0 | 6.5 | 3.5 |

Drum Bay

| Date | 3/25 | 4/16 | 5/15 | 6/19 | 7/10 | 7/30 | 8/28 | 9/17 | 10/8 | 10/29 |
|------------|---------|---------|---------|---------|-------|-------|---------|---------|------|-------|
| Water type | surface | surface | surface | surface | mixed | mixed | surface | surface | soil | soil |
| Salinity | 8.0 | 6.0 | 5.0 | 3.0 | 15.0 | 17.0 | 7.5 | 6.5 | 9.0 | 7.5 |

the year. Spartina alterniflora shares sub-dominance with S. patens. Juncus roemerianus is also present but only as a minor constituent.

When the PSS is between 15 and 25 ppt as in Dead Duck Pass, Juncus roemerianus is the dominant species. Spartina patens is then relegated to a very minor role. Spartina alterniflora is quite important as the chief species found growing between the clumps of Juncus roemerianus. Distichlis spicata plays only a minor role in the composition of this community.

In those areas with the highest PSS and regular flooding, such as Drum Bay, an almost pure community of Spartina alterniflora is found.

Seasonality

Figures 9-14 show the seasonal variation in species composition over time for the mixed sites.

Figures 9 and 10 indicate that Distichlis spicata maintains dominance throughout the year in Petain Lagoon. A slight increase in the importance of Spartina patens is exhibited in the months of November and December. Correspondingly, a decrease in Distichlis spicata and Spartina cynosuroides is seen.

Figures 11 and 12 indicate a period of high importance for Distichlis spicata in early spring with a decline to a low in early winter. This change at Bottle Lagoon is similar to the change at Petain Lagoon. Correspondingly there is an increase from early spring

toward winter in the ~~importance~~ of both Spartina alterniflora and Spartina patens.

Figures 13 and 14 for Dead Duck Pass show that Juncus roemerianus remains relatively constant in its importance throughout the year. Only Spartina alterniflora seems to display any change in its importance during the year, with a slight increase at the time of its flowering. Spartina patens and Distichlis spicata show an inconspicuous level of importance at all times.

Conclusions

From the work completed it appears that spring to early summer (weeks 18-34 of the year; April - mid-July) is the best period of time to categorize the communities. It is during this time of the year that the communities appear most stable in regard to species composition. This allows a strong correlation to be drawn between the salinity of the region (PSS) and the dominant species of the community. As such, this would seem to be the best period in which to sample the marsh via air or land for differences in vegetation and salinity.

A slight discrepancy in the sampling methods was noticed. Apparently the line transect method underestimates the smaller stemmed species such as Spartina patens. This is probably due to the difficulty of discerning individual plants. In conclusion, both methods are valid and when used together provide a good measure of marsh vegetation composition.




Figure 9. Clipped importance values from Petain Lagoon plotted against weeks of the year 1975.

- *Distichlis spicata*
- ▼ *Spartina cynosuroides*
- *Spartina patens*




Figure 10. Line transect importance values from Petain Lagoon plotted against weeks of the year 1975.

- *Distichlis spicata*
- ▼ *Spartina cynosuroides*
- *Spartina patens*




Figure 11. Clipped importance values from Bottle Lagoon plotted against weeks of the year 1975.

- Distichlis spicata
- Juncus roemerianus
- ▼ Spartina alterniflora
- Spartina patens




Figure 12. Line transect importance values from Bottle Lagoon plotted against weeks of the year 1975.

- Distichlis spicata
- Juncus roemerianus
- ▼ Spartina alterniflora
- Spartina patens




Figure 13. Clipped importance values from Dead Duck Pass plotted against weeks of the years 1975 and early 1976.

- Distichlis spicata
- Juncus roemerianus
- ▼ Spartina alterniflora
- Spartina patens




Figure 14. Line transect importance values from Dead Duck Pass plotted against weeks of the years 1975 and early 1976.

- Distichlis spicata
- Juncus roemerianus
- ▼ Spartina alterniflora
- Spartina patens

LITERATURE CITED

- Adams, D.A. 1963. Factors influencing vascular plant zonation in North Carolina salt marshes. *Ecol.* 44(3): 445-456.
- Albrecht, W.A. 1941. Soil organic matter and in availability for plants. *Soil Sci.* 51: 487-494.
- Allen, P.F. 1950. Ecological basis for land use planning in Gulf Coast marsh lands. *J. Soil Wat. Conserv.* 5(2): 57-62.
- Aurand, D. 1968. The seasonal and spatial distribution of nitrate and nitrite in the surface waters of two Delaware salt marshes. M.S. Thesis. Univ. of Dela.
- Babcock, K.M. 1967. The Influence of Water Depth and Salinity on Wiregrass and Saltmarsh grass. La. Wild. and Fisheries Commission.
- Blum, J.L. 1968. Salt marsh Spartinas and associated algae. *Ecol. Monogr.* 38: 199-222.
- _____. 1969. Nutrient changes in water flooding the high salt marsh. *Hydrobiologia* 34: 95-99.
- Bouleau, P.F. and D.A. Adams. 1956. Factors in vegetational zonation of salt marshes near Southport, North Carolina. *Bull. Eco. Am.* 37(3): 68.
- Bourn, W.S. and C. Cottam. 1950. Some biological effects of ditching tidewater marshes. U.S. Dept. Int. Rept. 1.
- Bray, J.R. 1962. Estimates of energy budgets for a Typha (cattail) marsh. *Science.* 136: 1119-1120.

- Buttery, B.R., W.T. Williams and J.M. Lambert. 1965. Competition between Glyceria maxima and Phragmites communis in the region of Surlingham Broad. II. The Fen Gradient. J. Ecol. 53: 183-195.
- Canfield, R. 1941. Application of the live interception method in sampling range vegetation. J. Forestry. 39: 388-394.
- Chabreck, R. 1970. Marsh zones and vegetative types in the Louisiana Coastal Marshes. Ph.D. Thesis. L.S.U.
- _____, T. Joanen and A.W. Palmisano. 1968. Vegetation type map of the Louisiana coastal marshes. Louisiana Wildlife and Fish. Comm.
- _____ and A.W. Palmisano. 1973. The effects of Hurricane Camille on the marshes of the Mississippi River. Ecol. 54: 1118-1123.
- Chamberlain, J.L. 1957. An ecological study of a gulf coast marsh, Rockefeller Refuge, Grand Chenier, La. Unpubl. PhD. Diss. Univ. of Tenn.
- _____. 1959. Influence of Hurricane Audrey on the Coastal Marsh of Southwestern Louisiana. La. State Univ. Coastal Stud. Inst., Tech. Rep. No. 10, Part B.
- Chapman, V.J. 1960. Salt Marshes and Salt Deserts of the World. Leonard Hill (Books) Ltd. London 392 pp.
- _____. 1940a. Succession on the New England salt marshes. Ecol. 21: 279-282.
- _____. 1940b. Studies in salt marsh ecology. Section VI and VII. Comparison with marshes on the east coast of North America. J. Ecol. 28: 118-152.

- Childress, U.R. 1960. Oyster and Fisheries Investigations of San Antonio and Espiritu Santo Bay (area m-5). Job report No. C-2, C-3. Texas Game and Fish Comm., Marine Div.
- _____. 1962. An Ecological Survey of Upper Galveston and Trinity Bays. Texas Game and Fish Commission Project Report No. M-2-R-3.
- Cibula, W.G. 1972. Application of Remotely Sensed Multispectral Data to Automated Analysis of Marshland Vegetation. Inference to the Location of Breeding Habits of the Salt Marsh Mosquito (Aedes sollicitans) National Aeronautics and Space Administration-Earth Resources Laboratory. Report No. 020.
- Coleman, J.M. 1966. Recent Coastal Sedimentation: Central Louisiana Coast. Louisiana State University Press. Baton Rouge, La.: 1-73.
- Conrad, H.S. 1935. The plant associations of central Long Island. Am. Midland Nat. 16(4): 433-516.
- Cox, G.W. 1972. Laboratory manual of general ecology. Wm. C. Brown Co., Dubuque, Iowa. 195 pp.
- Curl, H. 1959. The Hydrography of the Inshore Northeastern Gulf of Mexico. Publs. Inst. Mar. Sci. Univ. Texas 6: 193-205.
- de la Cruz, A.A. 1974. Primary productivity of coastal marshes in Mississippi. Gulf Research Reports 4(3): 351-356.
- _____. 1973. The role of tidal marshes in the productivity of coastal waters. Assoc. S.E. Biol. Bull. 20: 147-156.
- _____ and B.C. Gabriel. 1973. Caloric, elemental, and nutritive changes in decomposing Juncus roemerianus leaves. Ecol. 55: 882-886.

- _____. 1965. A study of particulate organic Detritus in a Georgia salt marsh-estuarine ecosystem. PhD. Diss. Univ. of Georgia, Athens, 110 pp.
- Davis, J.H. 1943. The natural features of southern Florida, especially the vegetation and the Everglades. Florida Geol. Survey Geol. Bull.
- Day, D.S. 1959. General Ecological Survey of the Matagorda Bay area. Job. report No. C-2. Texas Fish and Game Comm., Marine Div.
- Eggler, W.A. 1961. Vegetation of the drainage basin of Grand Bayou Blue. In R.J. Russell (Ed.), Louisiana Coastal Marsh Ecology. Tech. Rep. No. 14. Coastal Studies Inst., La. State Univ., Baton Rouge.
- Eichorn, R.E. and C.J. Guice. 1969. Louisiana coastal area marsh vegetation types. Bureau of Sport Fisheries and Wildlife, 4 p. memo.
- Eleuterius, L.N. 1973. The Marshes of Mississippi. In Cooperative Gulf of Mexico Estuarine Inventory and Study, Mississippi. Gulf Coast Research Laboratory (publisher). J.Y. Christmas-director and editor.
- _____. 1971. Recent changes in the Louisiana marsh near Vermilion Bay. Gulf Research Reports 3(2): 259-263
- Ensminger, A.B. and L.G. Nichols. 1957. Hurricane damage to Rockefeller Refuge. Proc. SE Assoc. Game Fish. Comm. 11: 52-56.
- Flowers, M. 1973. Vegetational zonation in two successional brackish marshes of the Chesapeake Science 14(3): 197-200.

- Fogarty, M.J. 1965. Biological studies within the coastal marsh area of Lake Calcasieu, La. Unpubl. Master's Thesis, L.S.U., Baton Rouge.
- Foster, W.A. 1968. Studies on the distribution and growth of Juncus roemerianus in southeastern Brunswick County, North Carolina, M.S. Thesis. Univ. of Michigan.
- Gabriel, B.C. and A.A. de la Cruz. 1974. Species composition, standing stock, and net primary production of a salt marsh community in Mississippi. Chesapeake Science 15(2): 72-77.
- Ganong, W.F. 1903. Vegetation of the Bay of Fundy marshes. Bot. Gaz. 36: 349-367.
- Good, R.E. 1965. Salt marsh vegetation, Cape May, New Jersey. The Bulletin-New Jersey Acad. of Sci. 10(1): 1-1.
- Gorham, E. 1953. Chemical studies on the soils and vegetation of water-logged habitats in the English Lake District. J. Ecol. 41: 345-360.
- _____ and W.H. Pearsall. 1956. Production ecology. III. Shoot production in Phragmites in relation to habitat. Oikos 7: 206-214.
- Gould, E. and J. Ewan. 1975. Phytogeographic and ecologic relationships of the flora of Breton Island, Louisiana. Tulane Studies in Zoology and Botany. 19(1, 2): 26-36.
- Griffitts, T.H.D. 1928. Some phases of the salt marsh mosquito problem in the South Atlantic and Gulf States. Proc. N.J. Mosquito Ext. Assoc. 87-91.

_____. 1929. Salt marsh vegetation in relation to salt marsh mosquito breeding in the South Atlantic and Gulf States.

Proc. N.J. Mosquito Ext. Assoc. 108-116.

Harris, V.T. and R.H. Chabreck. 1958. Some effects of Hurricane Audrey to the marsh at Marsh Island, Louisiana. Proc. La. Acad. Sci. 21: 47-50.

Harper, R.M. 1918. Some dynamic studies of Long Island vegetation. Plant World. 21: 38-46.

Harshberger, J.W. 1900. An ecological study of the New Jersey strand flora. Proc. Acad. Nat. Sci., Phil., 1900: 623-271.

_____. 1909. The vegetation of the salt marshes and of the salt and fresh water ponds of northern coastal New Jersey. Proc. Acad. Nat. Sci. Phil., 1909: 373-400.

Harshberger, J.W. 1911. An hydrometric investigation of the influences of sea water on the distribution of salt marshes and estuarine plants. Proc. Amer. Phil. Soc. 50: 457-484.

_____. 1916. The origin and vegetation of salt marsh pools. Proc. of Am. Phil. Soc. 55: 481-484.

Hoese, H.D. 1967. Effect of higher than normal salinities on salt marshes. Contr. In Marine Sci. 12: 249-261.

Hitchcock, E.W. 1972. Fragile nurseries of the sea-Can we save our salt marshes? National Geographic 141: 729-765.

Hinde, H.P. 1954. The vertical distribution of salt marsh phanerogams in relation to tide level. Ecol. Monogr. 24: 209-225.

Heald, E.J. 1969. The production of organic detritus in a South Florida estuary. PhD Diss. Univ. of Miami, 110 pp.

- Jackson, C.R. 1952. Some topographic and edaphic factors affecting distribution in a tidal marsh. Quart. J. Florida Acad. Sci. 15: 137-146.
- Jervis, R.A. 1964. Primary production in a freshwater marsh ecosystem. PhD Thesis, Rutgers Univ.
- Johnson, M. 1970. Preliminary report on species composition, chemical composition, biomass, and production of marsh vegetation in the upper Patuxent estuary, Maryland. In final technical report for the period August 1968 to August 1970. Office of Water Resources. Natural Resources Inst., Univ. of Maryland: NRI Ref. No. 71-6.
- Johnson, D.S. and H.H. York. 1915. The relation of plants to tide levels, a study of factors affecting the distribution of marine plants at Cold Spring Harbor, Long Island, N.Y. Carnegie Inst. Wash. Publ. 206: 161 pp.
- Juneau, C.L., Jr. 1975. An Inventory and Study of the Vermilion Bay-Atchafalaya Bay Complex. Technical Bulletin No. 13. Louisiana Wildlife and Fisheries Commission Oysters, Water Bottoms and Seafoods Division.
- Kearney, T.H., Jr. 1900. The plant covering of Ocracoke Island. A study in the ecology of the North Carolina strand vegetation. Contr. U.S. Nat. Herb. 5(5): 261-319.
- _____. 1901. Report on a botanical survey of the Dismal Swamp region. Contr. U.S. Nat. Herb. 5(6): 321-550.
- Keefe, C.S. 1972. Marsh production: a summary of the literature. Contr. in Marine Sci. 16: 163-181.

Kerwin, J.A. 1966. Classification and structure of the tidal marshes of Potomac River, Virginia. ASB Bulletin 13: 40. Abstract.

_____ and R.A. Pedigo. 1971. Synecology of a Virginia salt marsh. Chesapeake Science 12(3): 125-130.

Kirby, C.J., Jr. 1971. The annual net primary production and decomposition of the salt marsh grass Spartina alterniflora Loisel in the Barataria Bay Estuary, Louisiana. PhD Diss. L.S.U., Baton Rouge.

Kurz, H.K. and K. Wagner. 1957. Tidal marshes of the Gulf and Atlantic coasts of northern Florida and Charleston, S.C. Florida St. Univ. Stud. 24: 1-168.

Latter, P.M. and J.B. Cragg. 1967. The decomposition of Juncus squarrosus leaves and microbiological changes in the profile of Juncus moor. J. Ecol. 55: 465-481.

Lemaire, R.J. 1961. A preliminary annotated checklist of the vascular plants of the marshes and included higher lands of St. Bernard Parish, La. Proc. La. Acad. of Sci. 24: 56-70.

Linton, T.L. 1968. A description of the south Atlantic and Gulf Coast marshes and estuaries. Proc. Marsh and Est. Manag. Symp. L.S.U., Baton Rouge.

Lloyd, F.E. and S.M. Tracey. 1901. The insular flora of Mississippi and Louisiana. Bull. Torrey Bot. Club. 28: 61-101.

Lowe, E.N. 1921. Plants of Mississippi. Miss. State Geol. Survey Bull. 17.

- Martin, A.C., N. Hotchkiss, F.M. Uhler and W.S. Bourn. 1953.
Classification of wetlands of the U.S. Spec. Sci. Report
Wildlife No. 20. U.S. Dept. of the Int. 1-14.
- McCormick, J., R.R. Grant, Jr., and R. Patrick. 1970. Two
Studies of Tinicum Marsh. The Conservation Foundation.
- McIntyre, G.A. 1953. Estimation of plant density using live
transects. J. Ecology 41: 319-330.
- McNaughton and L.L. Wolf. 1973. General Ecology. Holt, Rine-
hart Winston, Inc. 710 pp.
- Miller, W.R. and F.E. Egler. 1950. Vegetation of the Wequete-
quock-Pawcatuck tidal marshes, Connecticut. Ecol. Monog.
20: 143-172.
- Mobberley, D.G. 1956. Taxonomy and distribution of the genus
Spartina. Iowa State College Journal of Science 30(4):
471-574.
- Mohr, C. 1901. Plant Life of Alabama. Geol. Surv. Ala. 6: 11-
825.
- Mooring, M.T., A.W. Cooper and E.D. Seneca. 1971. Seed germina-
tion and evidence for height ectophenes in Spartina alterniflora
from North Carolina. Amer. J. Bot. 58: 48-55.
- Morgan, M.H. 1961. Annual angiosperm production on a salt marsh.
M.S. Thesis, Univ. of Delaware, 34 pp.
- National Shoreline Study. 1971. Inventory Report-Lower Mississippi
Region. Louisiana Dept. of the Army. New Orleans District,
Corps of Engineers.

- Nichols, L.G. 1959. Geology of Rockefeller Wildlife Refuge and Game Preserve. Tech. Bull. La. Wildl. and Fisheries Comm. 35 pp.
- Niering, W.A. 1961. Tidal marshes, their use in scientific research. Conn. Arbor. Bull. 12: 3-7.
- Nixon, S.W. and C.A. Oviatt. 1973. Ecology of a New England salt marsh. Ecol. Monog. 43:463-498.
- Odum, E.P. and A.A. de la Cruz. 1963. Detritus as a major component of ecosystems. A.I.B.S. Bull. 13(3): 39-40.
- _____ and M.E. Fanning. 1973. Comparison of the productivity of Spartina alterniflora and Spartina cynosuroides in Georgia coastal marshes. Bull. Ga. Acad. Sci., 31: 1-12.
- O'Neil, T. 1949. The Muskrat in the Louisiana Coastal Marshes. La. Dept. of Wild Life and Fisheries. 152 pp.
- Palmer, E.L. 1941. Marshes and their environs. Nature Mag. 34: 205-212.
- Palmisano, A., Jr. 1970. Plant community-soil relationships in Louisiana coastal marshes. PhD Thesis. L.S.U., Baton Rouge.
- Penfound, W.T. 1952. Southern swamps and marshes. Bot. Rev. 18(6): 413-436.
- _____ and E.S. Hathaway. 1938. Plant communities in the marshlands of southeastern Louisiana. Ecol. Monog. 8: 1-56.
- _____ and M.E. O'Neal. 1934. The vegetation of Cat Island, Mississippi. Ecol. 15: 1-16.
- Pomeroy, L.R., R.E. Johannes, E.P. Odum and B. Roffman. 1969. The phosphorus and zinc cycles and productivity of the salt marsh. In. Proc. 2nd Natl. Symp. on Radioecology. D.J. Nilson and F.C. Evans editors: 414-419.

- Pullen, E.J. 1960. A study of the marsh and marine plants in upper Galveston and Trinity Bays. Texas Game and Fish Commission Marine Laboratory. Project Reports.
- Rankin, W.G. 1936. Lower Mississippi River Delta-Reports on the Geology of Plaquemines and St. Bernard Parishes. Geological Bulletin No. 8. Dept. of Conservation, Louisiana Geological Survey.
- Ranwell, D.S. 1964a. Spartina salt marshes in southern England. II. Rate and seasonal pattern of sediment. J. Ecol. 52: 79-94.
- _____. 1964b. Spartina salt marshes in southern England. III. Rates of establishment, succession and nutrient supply at Bridgwater Bay, Somerset. J. Ecol. 52: 95-105.
- Redfield, A.C. 1972. Development of a New England salt marsh. Ecol. Monog. 42: 201-237.
- Reed, J.F. 1947. The relation of the *Spartinetum glabra* near Beaufort, N.C. to certain edaphic factors. Amer. Midl. Nat. 38: 605-614.
- Reese, W.D. 1961. An annotated checklist of the plants of the Redfish Point Field Station of the University of Southwestern Louisiana. University of Southwestern La. 9 p. mimeo.
- Reimold, R.J. 1965. An evaluation of inorganic phosphate concentrations of Canary Creek marsh. M.S. Thesis. Univ. of Dela.
- _____ and F.C. Daiber. 1970. Dissolved phosphorus concentrations in a natural salt marsh of Delaware. Hydrobiologia. 36: 361-371.

- Robinson, W.O. 1930. Some chemical phases of submerged soil conditions. Soil Sci. 30: 197-217.
- Rounsefell, G.A. 1964. Preconstruction study of the fisheries of estuarine areas traversed by the Mississippi River-Gulf Outlet Project. Fishery Bull. Wildl. Ser. Us. 63: 373-393.
- Rudolphs, W. 1926. Investigations of mosquito problems carried on at the N.J. Agr. Exp. Sta. during the past year. Proc. N.J. Mosquito Est. A-soc. 33-51.
- Russell, R.J. 1936. Physiography of Lower Mississippi River Delta. La. Dept. of Cons. Geol. Survey Bull. No. 8: 1-199.
- Schelske, C.L. and E.P. Odum. 1961. Mechanisms maintaining high productivity in Georgia estuaries. Proc. Gulf Caribb. Fish Inst. 14: 75-80.
- Schultz, R.L. 1960. A survey of the macro-vegetation present in Aransas, Capano, Redfish, Port and St. Charles Bays, January 1, 1960 to December 31, 1960. Texas Game and Fish Comm. Lab. Project Reports.
- Shea, M.L., R.S. Warren and W.A. Niering. 1975. Biochemical and transplantation studies of the growth form of Spartina alterniflora on Connecticut salt marshes. Ecol. 56: 461-466.
- Shiflet, T.N. 1963. Major ecological factors controlling plant communities in Louisiana marshes. J. Range Mgt. 16 (5): 231-234.
- Smalley, A.E. 1959. The growth cycle of Spartina and its relation to the insect populations in the marsh. Proc. Salt Marsh Conf., Mar. Inst. Univ. of Ga., Sapelo Island, Ga. 96-100.

- Smith, W.G. 1970. Spartina "die back" in Louisiana marshlands. Coastal Studies Bull. No. 5. Special Sea Grant Issue.
- St. Amant, L.S. 1959. Louisiana Wildlife Inventory. La. Wildl. and Fisheries Comm. New Orleans, 329 pp.
- Strong, C.W. 1966. An improved method of obtaining density from line transect data. Ecology 47: 311-313.
- Stroud, L.M. and A.W. Cooper. 1968. Color infrared aerial photographic interpretation and net primary productivity of regularly flooded North Carolina salt marsh. Univ. of North Carolina Water Resources Inst., Report No. 14.
- Taylor, N. 1939. Salt tolerance of Long Island salt marsh plants. N.Y. State Museum Cir. 23.
- Teal, J.M. 1962. Energy flow in the salt marsh ecosystem of Georgia. Ecol. 43(4): 614-624.
- _____ and M. Teal. 1969. Life and Death of the Salt Marsh. Audubon/Ballantine. 274 pp.
- Tharp, B.C. 1926. Structure of Texas vegetation east of the 98th meridian. Univ. Texas Bull. 2606.
- Thorne, R.F. 1954. Flowering plants of the waters and shores of the Gulf of Mexico. In Marmer, H.A. (ed.), Gulf of Mexico, Its Origins, Waters and Marine Life. U.S. Fish and Wildlife Serv. Fishery Bull. No. 89.
- Tidwell, Mac A. 1973. The Tabanidae (Diptera) of Louisiana. Tulane Studies in Zoology and Botany. 18(1 & 2): 1-95.

- Uhler, F.M. and N. Hotchkiss. 1968. Vegetation and its succession in marshes and estuaries along the south Atlantic and Gulf Coasts. Proc. Marsh and Estuaries Management Symp. L.S.U., Baton Rouge.
- Uphof, J.C. 1941. Halophytes. Bot. Rev. 7: 1-58.
- Viosca, P. 1928. Louisiana wetlands and the value of their wildlife and fishery resources. Ecol. 9: 216-230.
- _____. 1933. Louisiana Out-ofdoors. 1-187.
- Waits, E.D. 1967. Net primary productivity of an irregularly flooded North Carolina salt marsh. PhD Diss., North Carolina State Univ., 113 pp.
- Walker, E.R. 1939. Biotic succession in a coastal salt marsh. Proc. of the Oklahoma Acad. of Sci. for 1939. 95-97.
- Wells, B.W. 1928. Plant communities of the coastal plain of North Carolina and their successional relations. Ecol. 9: 230-242.
- Williams, R.B. and M.B. Murdoch. 1972. Compartmental analysis of the production of Juncus roemerianus in a North Carolina salt marsh. Chesapeake Sci. 13: 69-79.
- Woodhouse, W.W., Jr., E.D. Seneca, and S.W. Broome. 1974. Propagation of Spartina alterniflora for Substrate Stabilization and Salt Marsh Development. Tech. Mem. No. 46, U.S. Army Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, Va., 22060.